

Research provides new insights into antibiotics and pig feeds

January 16 2012



Antibiotics in pig feed are increasing antibiotic resistant genes in pigs. Photo courtesy of MSU.

Antibiotics in pig feed increased the number of antibiotic resistant genes in gastrointestinal microbes in pigs, according to a study conducted by Michigan State University and the U.S. Department of Agriculture's Agricultural Research Service.

Published in the current edition of the *Proceedings of the National Academy of Sciences*, the comprehensive study focused on understanding

the effects of conventional, in-feed [antibiotics](#) in U.S. farms.

For decades, many producers of pigs, chickens and other farm animals have used antibiotics not only to protect their livestock from disease, but also to boost growth rates and enhance feed efficiency, a measure of how well animals convert feed into weight gains.

Scientists don't know precisely how antibiotics enhance growth rates and feed efficiency, but they are concerned that on-farm use of these medications may contribute to the development of strains of microbes resistant to conventional antibiotics, which are potentially harmful to humans and animals, said James Tiedje, MSU University Distinguished Professor of microbiology and molecular genetics and of crop and soil sciences.

“The growth of antibiotic resistance in pathogens is a huge challenge for society around the world,” said Tiedje, an MSU AgBioResearch scientist. “Studies to understand what contributes to the spread and what interventions can help control the problem are vital.”

Additional findings include:

- Both diversity and abundance of antibiotic resistance [genes](#) increased in the intestinal microbial communities of the pigs treated with antibiotics. Longer term studies are needed.
- Some of the genes found in the treated pigs were unexpected and usually linked to antibiotics not used in the study.
- Microbial genes associated with production and use of energy by [microbes](#) increased in abundance in the antibiotic-fed pigs, which may shed light on how antibiotics increase livestock growth and feed efficiency.

- E. coli populations increased in the intestines of the treated [pigs](#). Further study is needed to clarify this observation.

“To our knowledge, this study is the first of its kind to look at the collateral impacts of in-feed antibiotic use in farm animals, using a comprehensive approach to detect shifts in the function and the makeup or membership of the microbial community in the model animal’s gastrointestinal tract,” said Torey Looft, USDA researcher.

Additional MSU researchers included Tim Johnson, doctoral student; Robert Stedtfeld, civil and environmental engineering research associate; Woo Jun Sul, doctoral student; Tiffany Stedtfeld, civil and environmental engineering technical aide; Benli Chai, information technologist, Center for Microbial Ecology; James Cole, assistant professor at the Center for Microbial Ecology; and Syed Hashsham, civil and environmental engineering professor.

More information: In-feed antibiotic effects on the swine intestinal microbiome, Torey Looft et al., *PNAS*, [doi:10.1073/pnas.1120238109](https://doi.org/10.1073/pnas.1120238109)

Provided by Michigan State University

Citation: Research provides new insights into antibiotics and pig feeds (2012, January 16) retrieved 17 July 2024 from <https://phys.org/news/2012-01-insights-antibiotics-pig.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.