


# Not all who wander are lost

January 9 2012



Measurements of the metal content of stars in the disk of our Galaxy, using stars observed by SDSS-III's SEGUE-2 survey. Horizontal lines describe where SEGUE data measure the chemical composition of stars near and above the plane of the disk. The bottom panel shows the decrease in metal content as the distance from the Galactic center increases for stars near the plane of the Milky Way disk. In contrast, the metal content for stars far above the plane, shown in the upper panel, is nearly constant at all distances from the center of the Galaxy. The image of the Milky Way is from the Two-Micron All Sky Survey. Image Credit: Judy Cheng and Connie Rockosi (University of California, Santa Cruz) and the 2MASS Survey

(PhysOrg.com) -- Some stars have orbits that take them to interesting places, and they have interesting stories to tell about how they were formed. For more than a decade, the Sloan Digital Sky Survey (SDSS) has been mapping the stars in our galaxy. At today's meeting of the American Astronomical Society, astronomers Judy Cheng and and

Connie Rockosi (University of California, Santa Cruz) presented new evidence that will help answer long-standing questions about the history of disk of our galaxy.

These SDSS results come from the Sloan Extension for Galactic Understanding and Exploration 2 (SEGUE-2) in the SDSS-III project, which has measured the motions and chemical compositions of more than 118,000 stars. Some of those stars are in the disk of our galaxy, which is what we see as the bright band of stars in the night sky that we call the Milky Way. Stars in the disk [orbit](#) around the center of the galaxy. Most of the orbits line up in a flat plane like planets around the Sun — but a few orbit to the beat of a different drummer.

"Some disk stars have orbits that take them far above and below the plane of the Milky Way," explains Connie Rockosi, an astronomer at University of California Santa Cruz, and principal investigator for the SEGUE-2 survey. "We want to understand what kinds of stars those are, where they came from, and how they got there."

The orbits of these stars make them clearly different from mainstream Milky Way stars — but new research shows that their chemical composition also makes them unique. A team led by Judy Cheng of University of California Santa Cruz studied the amount of metals in such stars at different locations in the galaxy.

By making these measurements, Cheng's team was able to look back in time at how the Milky Way disk grew. The first generation of stars consisted entirely of hydrogen and helium. Over time, those early stars turned some of their hydrogen and helium into [heavier elements](#), like calcium or iron — and when those stars died, the heavier elements they produced became part of the next generation of stars. As new stars were born and the Milky Way disk grew, each generation had more calcium, iron, and other heavy elements. Thus, we can learn which parts of our

galaxy have seen several generations of stars come and go, simply by looking at the metal content of stars in that part of the galaxy.

Cheng's team surveyed the metal content of thousands of stars in the disk. Near the plane of the galactic disk — home of the ordinary disk stars — stars closer to the center of the galaxy have higher metal content than those farther from the galactic center. "That tells us that the outer disk of our Galaxy has formed fewer generations of stars than the inner disk — meaning that the Milky Way disk grew from the inside out," says Cheng.

But then Cheng studied the "different drummer" stars, those that are clearly part of the Milky Way disk but show up far above or below the disk plane. She found that the amount of heavier elements in those stars doesn't follow the same trend — everywhere she looked in that part of the galaxy, stars had low metal content. "The fact that the metal content of those stars is the same everywhere is a new piece of evidence that can help us figure out how they got to be so far away from the plane," Rockosi says.

What we do not yet know is whether these stars were born with such strange orbits, or whether something happened in the past to put them on these unique paths. "If these stars were born with these orbits," says Cheng, "they were born at the same rate all over the galaxy. If they were born with regular orbits, then whatever happened to them must have been very efficient at mixing them up and erasing any patterns in the metal content, such as the inside-out trend we see in the plane." Possible explanations for such efficient mixing include long-ago collisions between our galaxy and its neighbors, or the effect of spiral arms sweeping through the disk. Cheng's observations will help determine whether such major events in the lives of these stars caused them to wander far from their birthplace.

Disk stars are observed far from the plane in many other [galaxies](#), so solving the puzzle presented by these [stars](#) observed by SDSS will help us understand a basic part of how spiral galaxies like the [Milky Way](#) form.

Provided by Sloan Digital Sky Survey

Citation: Not all who wander are lost (2012, January 9) retrieved 27 April 2024 from <https://phys.org/news/2012-01-lost.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.