

Alan Turing's 1950s tiger stripe theory proved

February 19 2012



Image: Wikipedia.

Researchers from King's College London have provided the first experimental evidence confirming a great British mathematician's theory of how biological patterns such as tiger stripes or leopard spots are formed.

The study, funded by the Medical Research Council and to be published online in [Nature Genetics](#), not only demonstrates a mechanism which is likely to be widely relevant in vertebrate development, but also provides confidence that chemicals called morphogens, which control these patterns, can be used in regenerative medicine to differentiate stem cells into tissue.

The findings provide evidence to support a theory first suggested in the

1950s by famous code-breaker and mathematician Alan Turing, whose centenary falls this year. He put forward the idea that regular [repeating patterns](#) in [biological systems](#) are generated by a pair of morphogens that work together as an 'activator' and 'inhibitor'.

To test the theory the researchers studied the development of the regularly spaced ridges found in the roof of the mouth in mice. Carrying out experiments in [mouse embryos](#), the team identified the pair of morphogens working together to influence where each ridge will be formed. These chemicals controlled each other's expression, activating and inhibiting production and therefore controlling the generation of the ridge pattern.

The researchers were able to identify the specific morphogens involved in this process – FGF (Fibroblast Growth Factor) and Shh (Sonic Hedgehog – so-called because laboratory fruit flies lacking the fly version have extra bristles on their bodies). They showed that when these morphogens' activity is increased or decreased, the pattern of the ridges in the mouth palate are affected in ways predicted by Turing's equations. For the first time the actual morphogens involved in this process have been identified and the team were able to see exactly the effects predicted by Turing's 60-year-old speculative theory.

Dr Jeremy Green from the Department of Craniofacial Development at King's Dental Institute said: 'Regularly spaced structures, from vertebrae and hair follicles to the stripes on a tiger or zebrafish, are a fundamental motif in biology. There are several theories about how patterns in nature are formed, but until now there was only circumstantial evidence for Turing's mechanism. Our study provides the first experimental identification of an activator-inhibitor system at work in the generation of stripes – in this case, in the ridges of the mouth palate.

'Although important in feeling and tasting food, ridges in the mouth are

not of great medical significance. However, they have proven extremely valuable here in validating an old theory of the activator-inhibitor model first put forward by Alan Turing in the 50s.

'Not only does this show us how patterns such as stripes are formed, but it provides confidence that these morphogens (chemicals) can be used in future regenerative medicine to regenerate structure and pattern when differentiating [stem cells](#) into other tissues.

'As this year marks Turing's centenary, it is a fitting tribute to this great mathematician and computer scientist that we should now be able to prove that his theory was right all along!'

Provided by King's College London

Citation: Alan Turing's 1950s tiger stripe theory proved (2012, February 19) retrieved 26 April 2024 from <https://phys.org/news/2012-02-alan-turing-1950s-tiger-stripe.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
