

Borexino Collaboration succeeds in spotting pep neutrinos emitted from the sun

February 9 2012, by Bob Yirka



View of the Borexino "Stainless Steel Sphere" (SSS) from from the "Water Tank"

(PhysOrg.com) -- To learn more about how the sun works, scientists study particles that are emitted from it into space due to thermonuclear reactions that occur inside; by applying known physics principles, they can then deduce which sort of nuclear reactions are taking place. As one example, researchers have been able to identify high energy proton to proton interactions that are described as *pp* neutrinos by detecting them when they reach Earth.

Now, researchers deep beneath the ground in a mountain in Italy, working together in a group known as the Borexino Collaboration, have spotted the more elusive proton to electron to proton neutrino, a *pep*

reaction that results in the formation of [deuterium](#), a heavy form of [hydrogen](#). To detect them, the team, as they describe in their paper published in [Physical Review Letters](#), the team had to develop a method of filtering out virtually all other neutrinos, including those from [outer space](#).

To detect the presence of neutrinos, researchers build underground facilities to use the Earth's natural filtering abilities to remove particle clutter. Then, they fill a big vat with a special kind of liquid that reacts with the type of neutrino they are looking for. When one of the neutrinos strikes the liquid, a tiny flash or sparkle occurs. By measuring the number of sparkles that occur over a period of time the researchers can describe the amount of such neutrinos that are emitted by the sun, which helps to more fully understand the [nuclear reactions](#) that occur inside of it.

Pp neutrinos have been easy to count, they are plentiful and [high energy](#), which makes it easy to detect them when hitting the liquid. *Pep*, neutrinos on the other hand are low energy and more elusive and up till now have been mostly a theoretical concept. To detect their presence the team had to devise a means of filtering out virtually all other cosmic particles and then use a liquid that causes a sparkle when struck by a particle that has just 1.44 mega-electron-volts of energy, the distinctive signature of the *pep* neutrino. The team succeeded on both counts and were able to detect 3.1 *pep* neutrino strikes per day, per 100 tons of liquid.

The new technique for cleaning and filtering out unwanted particles is ground breaking work and likely will be used by other scientists looking to measure other [particles](#) in other research efforts.

More information: First Evidence of *pep* Solar Neutrinos by Direct Detection in Borexino, *Phys. Rev. Lett.* 108, 051302 (2012).

[DOI:10.1103/PhysRevLett.108.051302](https://doi.org/10.1103/PhysRevLett.108.051302)

We observed, for the first time, solar neutrinos in the 1.0–1.5 MeV energy range. We determined the rate of pep solar neutrino interactions in Borexino to be $3.1 \pm 0.6_{\text{stat}} \pm 0.3_{\text{syst}}$ counts/(day·100 ton). Assuming the pep neutrino flux predicted by the standard solar model, we obtained a constraint on the CNO solar neutrino interaction rate of

Citation: Borexino Collaboration succeeds in spotting pep neutrinos emitted from the sun (2012, February 9) retrieved 18 April 2024 from <https://phys.org/news/2012-02-borexino-collaboration-pep-neutrinos-emitted.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.