

With climate change, today's '100-year floods' may happen every three to 20 years: research

February 13 2012, by Jennifer Chu



An atmospheric image of Hurricane Irene on the U.S. East Coast in August 2011. Image: NOAA

Last August, Hurricane Irene spun through the Caribbean and parts of the eastern United States, leaving widespread wreckage in its wake. The Category 3 storm whipped up water levels, generating storm surges that swept over seawalls and flooded seaside and inland communities. Many hurricane analysts suggested, based on the wide extent of flooding, that Irene was a “100-year event”: a storm that only comes around once in a century.

However, researchers from MIT and Princeton University have found that with [climate change](#), such storms could make landfall far more

frequently, causing powerful, devastating storm surges every three to 20 years. The group simulated tens of thousands of storms under different climate conditions, finding that today's "500-year floods" could, with climate change, occur once every 25 to 240 years. The researchers published their results in the current issue of *Nature Climate Change*.

MIT postdoc Ning Lin, lead author of the study, says knowing the frequency of storm surges may help urban and coastal planners design seawalls and other protective structures.

"When you design your buildings or dams or structures on the coast, you have to know how high your seawall has to be," Lin says. "You have to decide whether to build a seawall to prevent being flooded every 20 years."

Lin collaborated with Kerry Emanuel, the Cecil and Ida Green Professor of Atmospheric Science at MIT, as well as with Michael Oppenheimer and Erik Vanmarcke at Princeton. The group looked at the impact of climate change on storm surges, using New York City as a case study.

To simulate present and future storm activity in the region, the researchers combined four climate models with a specific hurricane model. The combined models generated 45,000 synthetic storms within a 200-kilometer radius of Battery Park, at the southern tip of Manhattan.

They studied each climate model under two scenarios: a "current climate" condition representing 1981 to 2000 and a "future climate" condition reflecting the years 2081 to 2100, a prediction based on the Intergovernmental Panel on Climate Change's projections of future moderate carbon dioxide output. While there was some variability among the models, the team generally found that the frequency of intense storms would increase due to climate change.

Once they simulated storms in the region, the researchers then simulated the resulting storm surges using three different models, including one used by the National Hurricane Center (NHC). In the days or hours before a hurricane hits land, the NHC uses a storm-surge model to predict the risk and extent of flooding from the impending storm. Such models, however, have not been used to evaluate multiple simulated storms under a scenario of climate change.

Again, the group compared results from multiple models: one from the NHC which simulates storm surges quickly, though coarsely; another model that generates more accurate storm surges, though less efficiently; and a model in between, developed by Lin and her colleagues, that estimates relatively accurate surge floods, relatively quickly.

Today, a “100-year storm” means a surge [flood](#) of about two meters, on average, in New York. Roughly every 500 years, the region experiences towering, three-meter-high surge floods. Both scenarios, Lin notes, would easily top Manhattan’s seawalls, which stand 1.5 meters high.

But with added greenhouse gas emissions, the models found that a two-meter surge flood would instead occur once every three to 20 years; a three-meter flood would occur every 25 to 240 years.

“The highest [surge flood] was 3.2 meters, and this happened in 1821,” Lin says. “That’s the highest water level observed in New York City’s history, which is like a present 500-year event.”

Carol Friedland, an assistant professor of construction management and industrial engineering at Louisiana State University, sees the group’s results as a useful tool to inform coastal design — particularly, she notes, as most buildings are designed with a 60- to 120-year “usable lifespan.”

“The physical damage and economic loss that result from storm surge

can be devastating to individuals, businesses, infrastructure and communities,” Friedland says. “For current coastal community planning and design projects, it is essential that the effects of climate change be included in storm-surge predictions.”

This story is republished courtesy of MIT News (web.mit.edu/newsoffice/), a popular site that covers news about MIT research, innovation and teaching.

Provided by Massachusetts Institute of Technology

Citation: With climate change, today's '100-year floods' may happen every three to 20 years: research (2012, February 13) retrieved 22 April 2024 from <https://phys.org/news/2012-02-climate-today-year-years.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.