

Economizing chemistry, atom by atom

February 3 2012




Figure 1: A scandium catalyst (green spheres, center) can catalyze the addition of a phenylsilane (orange structure) to anisole (blue structure) to give cleaner products, using fewer reagents, than achieved previously. Credit: 2012 Zhaomin Hou

In chemistry, downsizing can have positive attributes. Reducing the number of steps and reagents in synthetic reactions, for example, enables chemists to boost their productivity while reducing their environmental footprint. This type of ‘atom economy’ could soon improve, thanks to a new rare-earth metal catalyst developed by Zhaomin Hou and colleagues at the RIKEN Advanced Science Institute, Wako. Their catalyst makes it simpler to modify aromatic carbon–hydrogen (C–H) bonds with silicon-bearing silyl ligands—a reaction step critical to pharmaceutical and materials science manufacturers alike.

Silicon, which is less electronegative than carbon or hydrogen atoms, can significantly alter the electronic characteristics of an organic molecule. Replacing the hydrogen atoms of an aromatic C–H group with silyl groups has emerged as an important strategy in industrial-scale chemical synthesis because these substituents can tune molecular reactivity,

enabling construction of elaborate chemical frameworks.

[Chemists](#) normally use transition metals such as platinum or rhodium to catalyze aromatic silylation reactions. But to achieve high conversions, these catalysts need to be mixed with additional hydrogen acceptor [reagents](#), which can generate unwanted waste products, including alkanes.

Hou and colleagues have pioneered studies into rare-earth metals, such as scandium, which have different catalytic properties to transition metals. Recently, they found that ‘half-sandwich’ scandium complexes, bonded on one side by a flat organic ring, showed unique activity and selectivity in the presence of carbon double bonds. This made investigations of unsaturated aromatic molecules a natural next step.

When the researchers mixed a methoxy–benzene compound called anisole with the half-sandwich scandium catalyst and a phenylsilane, they found that the silyl group substituted onto the aromatic ring with excellent selectivity and yields (Fig. 1). Furthermore, the catalyst did not require hydrogen acceptor reagents, and generated only H₂ gas as a by-product. Hou notes that this reaction is highly advantageous in terms of atom economy.

X-ray and spectroscopic measurements revealed that the working form of the catalyst, which contained a pair of ‘bridging’ [hydrogen atoms](#), activated the reaction by coordinating the anisole’s methoxy group to the rare-earth metal. According to Hou, this relatively strong interaction directs silylation to occur almost exclusively at the position adjacent to the methoxy unit on the aromatic ring—a ‘regioselectivity’ that outshines that of transition metal catalysts, whose weak oxygen–metal interactions often produce an undesirable mix of silylation isomers.

The team will continue to explore new approaches to improving catalytic

sustainability and selectivity by tapping into the extraordinary properties of rare-earth metals.

More information: Oyamada, J., et al. Scandium-catalyzed silylation of aromatic C–H bonds. [Angewandte Chemie International Edition](#) 50, 10720–10723 (2011).

Takimoto, M., et al. Scandium-catalyzed regio- and stereospecific methylalumination of silyloxy/alkoxy-substituted alkynes and alkenes. [Journal of the American Chemical Society](#) 131, 18266–18268 (2009).

Provided by RIKEN

Citation: Economizing chemistry, atom by atom (2012, February 3) retrieved 6 May 2024 from <https://phys.org/news/2012-02-economizing-chemistry-atom.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.