

New study shows no evidence of groundwater contamination from hydraulic fracturing

February 16 2012

Hydraulic fracturing of shale formations to extract natural gas has no direct connection to reports of groundwater contamination, based on evidence reviewed in a study released Thursday by the Energy Institute at The University of Texas at Austin.

The study, released at the annual meeting of the [American Association for the Advancement of Science](#) (AAAS) in Vancouver, British Columbia, found that many problems ascribed to hydraulic fracturing are related to processes common to all [oil and gas drilling](#) operations, such as casing failures or poor [cement](#) jobs.

University researchers also concluded that many reports of contamination can be traced to above-ground spills or other mishandling of [wastewater](#) produced from [shale gas](#) drilling, rather than from hydraulic fracturing per se, said Charles "Chip" Groat, an Energy Institute associate director who led the project.

"These problems are not unique to hydraulic fracturing," he said.

The research team examined evidence contained in reports of groundwater contamination attributed to hydraulic fracturing in three prominent shale plays — the Barnett Shale in North Texas; the Marcellus Shale in Pennsylvania, New York and portions of Appalachia; and the Haynesville Shale in western Louisiana and northeast Texas.

The report identifies regulations related to shale [gas development](#) and

evaluates individual states' capacity to enforce existing regulations. In addition, university researchers analyzed public perceptions of hydraulic fracturing, as derived from popular media, scientific literature and online surveys.

"Our goal was to provide policymakers a foundation for developing sensible regulations that ensure responsible shale gas development," Groat said. "What we've tried to do is separate fact from fiction."

Faculty members from across The University of Texas at Austin campus participated in the research, which the Energy Institute funded. The Environmental Defense Fund also assisted in developing the scope of work and methodology for the study.

Groat said researchers will supplement the study released Thursday with an examination of reports relating to atmospheric emissions and seismic activity attributed to hydraulic fracturing, which have emerged as significant issues of concern in recent months.

Hydraulic fracturing involves the high-pressure injection of water, sand and chemicals into a shale bed, which causes the rock to shatter, releasing natural gas. The practice has been in use for decades but has come under scrutiny in recent years from environmentalists and others who fear it poses a threat to public health.

Other findings from the Energy Institute study include:

- [Natural gas](#) found in water wells within some shale gas areas (e.g., Marcellus) can be traced to natural sources and probably was present before the onset of shale gas operations.
- Although some states have been proactive in overseeing shale gas development, most regulations were written before the

widespread use of hydraulic fracturing.

- Media coverage of hydraulic fracturing is decidedly negative, and few news reports mention scientific research related to the practice.
- Overall, surface spills of fracturing fluids pose greater risks to groundwater sources than from hydraulic fracturing itself.
- The lack of baseline studies in areas of shale gas development makes it difficult to evaluate the long-term, cumulative effects and risks associated with hydraulic fracturing.

Groat also provided an overview of two other Energy Institute initiatives related to the use of hydraulic fracturing in shale gas development.

The first project, which will commence in April, is a detailed case study focusing on claims of [groundwater contamination](#) in North Texas' Barnett Shale. The research will entail an examination of various aspects of shale gas development, including site preparation, drilling, production, and handling and disposal of flow-back water. Researchers also will identify and document activities unrelated to shale gas development that have resulted in water contamination. It will also assess the quantity of fresh groundwater used in shale gas development and evaluate ways to reduce the amount.

A second project, currently under development, would include a field and laboratory investigation of whether hydrological connectivity exists between water in the units above and below the shale unit being fractured as a result of the fracturing process. As envisioned, the project calls for university researchers to conduct field sampling of [hydraulic fracturing](#) fluid, flow-back water, produced water, and water from aquifers and other geologic units within the Barnett Shale.

Provided by The Energy Institute at UT Austin

Citation: New study shows no evidence of groundwater contamination from hydraulic fracturing (2012, February 16) retrieved 5 April 2024 from <https://phys.org/news/2012-02-evidence-groundwater-contamination-hydraulic-fracturing.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.