

Fossil teeth of *Gigantopithecus* found from Yunnan-Guizhou plateau

February 29 2012



Fig. 1: Fossil teeth of *Gigantopithecus blacki* from Baeryan cave, Bijie county, Guizhou Province. 1-occlusal view, 2-lateral view. Credit: ZHAO Lingxia

Pleistocene *Gigantopithecus blacki* is the largest species of all extinct and extant primates. Its diet, distribution and evolution remained unclear. According to a paper in press in the journal of *Quaternary International* (2012), paleoanthropologists from Institute of Vertebrate Paleontology and Paleoanthropology (IVPP), Chinese Academy of Sciences, reported fossil teeth of Early Pleistocene *Gigantopithecus blacki* and associated mammalian fauna from Baeryan Cave, Bijie County in Guizhou Province, providing new evidence for analysis on this giant ape's diet, living habitat, distribution and evolution.

The Baeryan cave fossil site located in the Yunnan-Guizhou Plateau, is

the highest site in altitude among known sites of *G. blacki*, and was exposed when a local farmer made excavations in the 1980s. Five isolated teeth of *Gigantopithecus blacki* and other mammalian fossils were quarried out in 2001-2004. All teeth crowns are well preserved. Compared with the large assemblage (IVPP laboratory, Beijing) of *Gigantopithecus* teeth from Juyuangdong cave, Liucheng County, Guangxi Autonomous Region, the morphology features of Baeryan *Gigantopithecus* teeth show no difference.

A total of 21 species were identified from the Baeryan Cave site, including *G. blacki*, *Ailuropoda microta*, *Pachycrocuta licneti*, *Cuon dubius*, *Ursus thibetanus primitinus*, *Panthera* sp., *Dicoryphocheirus ultimus*, *Muntiacus* sp., *Cervavitus fenqii*, *Cervus* cf. *unicolor*, *Budorcas* sp., *Caprinae* gen. et sp. indet., *Gazella* sp., *Leptobos* sp., *Tapirus sanyuanensis*, *Nestoritherium* sp., *Rhinoceros sinensis*, *Rhinopithecus* sp., *Stegodon orientalis*, *Rhizomys* sp., and *Hystrix subcristata*. They are the typical elements of the Early Pleistocene fauna in South China.




Fig. 2. Micro-CT section through mesial cusps of the upper second molar (GBB1-1), indicating an adaptation to feeding on hard objects, such as seeds and

nuts. Credit: ZHAO Lingxia

Enamel stable carbon isotope analysis on *Gigantopithecus* teeth indicates that it is a pure C₃ [diet](#) and lived in a forest habitat, which is clearly different from early hominins in South and East Africa, such as *Australopithecus africanus*, *Paranthropus robustus* and *Paranthropus boisei*, which had C₄ diets.

What kinds of C₃ food did *G. blacki* eat? The high frequency of caries indicates a diet rich in carbohydrates or starch. Fifteen carious teeth were identified from the total of 62 molars discovered in Longgudong cave. Thick enamel and blunt-cusp morphology of buccal [teeth](#) are related to the adaptation to feeding on hard objects, such as seeds and nuts. Both pits and scratches of microwear were commonly present on the occlusal surface under SEM, which implies an omnivorous diet. The microwear evidence and the morphological features suggest *G. blacki* had a frugivorous and highly fibrous diet. It was also likely *Gigantopithecus* fed on bamboo (C₃ plant) as part of its diet. More detailed analysis is necessary for further interpretation of the diet of the giant ape.




Fig. 3. Microwear by SEM showing the pits and scratches on the protoconid surface of a lower first molar of *Gigantopithecus blacki* from Juyuandong cave, implying an omnivorous diet. Credit: ZHAO Lingxia

The fossil records indicate that *Gigantopithecus* was distributed widely during the Early Pleistocene, not only in the Guangxi area of South China (such as Juyuandong cave in Liucheng, Mohui cave and Chuifeng cave in Tiandong, Sanhe cave in Chongzuo), but north to the Three Gorges Area of the Yangtze River (such as Longgupo site in Wushan and Longgudong cave in Jianshi), and also in the Guizhou Plateau, such as the Baeryan site in Bijie. In the Middle Pleistocene, *Gigantopithecus* withdrew southward to a limited area of South China, mainly in Guangxi Province (Heidong cave in Daxin, Bulalishan cave in Wuming, Nongmoshan cave in Bama), and more southward to Hainan island (Xinchong cave) and North Vietnam (Tham Khuyen [cave](#)). No fossil of Late Pleistocene *Gigantopithecus* has been recorded so far.

The climate and environmental change during the late Early Pleistocene might have had a strong effect on the habitat and diet of *Gigantopithecus*, and made it withdraw southward into the subtropical

forest where the climate was warm and wet. Homo had evolved and existed in South China since the Early Pleistocene. Enlarging Homo populations with increasing intelligence and abilities in tool-making and fire-controlling might apply competitive pressure on Gigantopithecus for territory and resources, and finally made Gigantopithecus go extinct through the middle to late Pleistocene.

This work was supported by the National Science Foundation of China, State Key Laboratory of Palaeobiology and Stratigraphy of Nanjing Institute of Geology and Palaeontology, and the Specific Basic Research Program of MOST of China.

More information: Zhao, L.X., Zhang, L.Z., New fossil evidence and diet analysis of Gigantopithecus blacki and its distribution and extinction in South China, Quaternary International (2012), [doi:10.1016/j.quaint.2011.12.016](https://doi.org/10.1016/j.quaint.2011.12.016)

Provided by Institute of Vertebrate Paleontology and Paleoanthropology

Citation: Fossil teeth of Gigantopithecus found from Yunnan-Guizhou plateau (2012, February 29) retrieved 28 April 2024 from <https://phys.org/news/2012-02-fossil-teeth-gigantopithecus-yunnan-guizhou-plateau.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--