


Lunar Reconnaissance Orbiter reveals recent geological activity on the Moon

February 21 2012




This shows the largest of the newly detected graben found in highlands of the lunar farside. The broadest graben is about 500 meters (1,640 feet) wide and topography derived from Lunar Reconnaissance Orbiter Camera (LROC) Narrow Angle Camera (NAC) stereo images indicates they are almost 20 meters

(almost 66 feet) deep. (Credit: NASA/Goddard/Arizona State University/Smithsonian Institution)

(PhysOrg.com) -- New images from NASA's Lunar Reconnaissance Orbiter (LRO) spacecraft show the moon's crust is being stretched, forming minute valleys in a few small areas on the lunar surface. Scientists propose this geologic activity occurred less than 50 million years ago, which is considered recent compared to the moon's age of more than 4.5 billion years.

A team of researchers analyzing high-resolution images obtained by the [Lunar Reconnaissance Orbiter](#) Camera (LROC) show small, narrow trenches typically much longer than they are wide. This indicates the lunar crust is being pulled apart at these locations. These linear [valleys](#), known as graben, form when the moon's crust stretches, breaks and drops down along two bounding faults. A handful of these graben systems have been found across the lunar surface.

"We think the moon is in a general state of global contraction because of cooling of a still hot interior," said Thomas Watters of the Center for Earth and Planetary Studies at the Smithsonian's National Air and Space Museum in Washington, and lead author of a paper on this research appearing in the March issue of the journal *Nature Geoscience*. "The graben tell us forces acting to shrink the moon were overcome in places by forces acting to pull it apart. This means the contractional forces shrinking the moon cannot be large, or the small graben might never form."



Graben are troughs formed when the lunar crust was stretched and pulled apart. This stretching causes the near-surface materials to break along two parallel normal faults, the terrain in between the twin faults drops down forming a valley. (Credit: Arizona State University/Smithsonian Institution)


The weak contraction suggests that the moon, unlike the terrestrial planets, did not completely melt in the very early stages of its evolution. Rather, observations support an alternative view that only the moon's exterior initially melted forming an ocean of molten rock.

In August 2010, the team used LROC images to identify physical signs of contraction on the lunar surface, in the form of lobe-shaped cliffs known as lobate scarps. The scarps are evidence the moon shrank globally in the geologically recent past and might still be shrinking today. The team saw these scarps widely distributed across the moon and concluded it was shrinking as the interior slowly cooled.

Based on the size of the scarps, it is estimated that the distance between the moon's center and its surface shrank by approximately 300 feet. The graben were an unexpected discovery and the images provide

contradictory evidence that the regions of the lunar crust are also being pulled apart.

"This pulling apart tells us the moon is still active," said Richard Vondrak, LRO Project Scientist at NASA's Goddard Space Flight Center in Greenbelt, Md. "LRO gives us a detailed look at that process."



Newly detected series of narrow linear troughs are known as graben, and they formed in highland materials on the lunar farside. Forces acting to pull the lunar crust apart formed the Virtanen graben, informally named for a nearby impact crater. These graben are located on a topographic rise with several hundred meters of relief revealed in topography derived from Lunar Reconnaissance Orbiter Camera (LROC) Narrow Angle Camera (NAC) stereo images (blues are lower elevations and reds are higher elevations). The rise is flanked by the rim of

a ~2.5 km diameter degraded crater. Photo by: NASA/GSFC/Arizona State University/Smithsonian Institution

As the LRO mission progresses and coverage increases, scientists will have a better picture of how common these young graben are and what other types of tectonic features are nearby. The graben systems the team finds may help scientists refine the state of stress in the lunar crust.

"It was a big surprise when I spotted graben in the far side highlands," said co-author Mark Robinson of the School of Earth and Space Exploration at Arizona State University, principal investigator of LROC. "I immediately targeted the area for high-resolution stereo images so we could create a three-dimensional view of the graben. It's exciting when you discover something totally unexpected and only about half the [lunar surface](#) has been imaged in high resolution. There is much more of the [moon](#) to be explored."

Provided by JPL/NASA

Citation: Lunar Reconnaissance Orbiter reveals recent geological activity on the Moon (2012, February 21) retrieved 23 April 2024 from <https://phys.org/news/2012-02-lunar-reconnaissance-orbiter-reveals-geological.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.