


Planck mission steps closer to the cosmic blueprint

February 13 2012



This all-sky image shows the distribution of carbon monoxide (CO), a molecule used by astronomers to trace molecular clouds across the sky, as seen by Planck (blue). A compilation of previous surveys (Dame et al. (2001)), which left large areas of the sky unobserved, has been superimposed for comparison (red). The outlines identify the portions of the sky covered by these surveys. Credits: ESA/Planck Collaboration; T. Dame et al., 2001

(PhysOrg.com) -- ESA's Planck mission has revealed that our Galaxy contains previously undiscovered islands of cold gas and a mysterious haze of microwaves. These results give scientists new treasure to mine and take them closer to revealing the blueprint of cosmic structure.

The new results are being presented this week at an international conference in Bologna, Italy, where [astronomers](#) from around the world are discussing the mission's intermediate results.


These results include the first map of [carbon monoxide](#) to cover the entire sky. Carbon monoxide is a constituent of the cold clouds that populate the Milky Way and other [galaxies](#). Predominantly made of hydrogen molecules, these clouds provide the reservoirs from which stars are born.

However, hydrogen molecules are difficult to detect because they do not readily emit radiation. Carbon monoxide forms under similar conditions and, even though it is much rarer, it emits light more readily and therefore is more easily detectable. So, astronomers use it to trace the clouds of hydrogen.

“Planck turns out to be an excellent detector of carbon monoxide across the entire sky,” says Planck collaborator Jonathan Aumont from the Institut d’Astrophysique Spatiale, Universite Paris XI, Orsay, France.

Surveys of carbon monoxide undertaken with radio telescopes on the ground are extremely time consuming, hence they are limited to portions of the sky where molecular clouds are already known or expected to exist.

“The great advantage of Planck is that it scans the whole sky, allowing us to detect concentrations of molecular gas where we didn’t expect to find them,” says Dr Aumont.



This all-sky image shows the distribution of carbon monoxide (CO), a molecule used by astronomers to trace molecular clouds across the sky, as seen by Planck. The inserts provide a zoomed-in view onto three individual regions on the sky where Planck has detected concentrations of CO: Cepheus, Taurus and Pegasus, respectively. Credits: ESA/Planck Collaboration

Planck has also detected a mysterious haze of [microwaves](#) that presently defies explanation.

It comes from the region surrounding the galactic centre and looks like a form of energy called synchrotron emission. This is produced when electrons pass through magnetic fields after having been accelerated by supernova explosions.

The curiosity is that the synchrotron emission associated with the galactic haze exhibits different characteristics from the synchrotron emission seen elsewhere in the Milky Way.

The galactic haze shows what astronomers call a 'harder' spectrum: its emission does not decline as rapidly with increasing energies.

Several explanations have been proposed for this unusual behaviour,

including higher supernova rates, galactic winds and even the annihilation of dark-matter particles.

So far, none of them has been confirmed and it remains puzzling.

“The results achieved thus far by Planck on the galactic haze and on the carbon monoxide distribution provide us with a fresh view on some interesting processes taking place in our Galaxy,” says Jan Tauber, ESA’s Project Scientist for Planck.

Planck’s primary goal is to observe the Cosmic Microwave Background (CMB), the relic radiation from the Big Bang, and to measure its encoded information about the constituents of the Universe and the origin of cosmic structure.

But it can only be reached once all sources of foreground emission, such as the galactic haze and the carbon monoxide signals, have been identified and removed.

“The lengthy and delicate task of foreground removal provides us with prime datasets that are shedding new light on hot topics in galactic and extragalactic astronomy alike,” says Dr Tauber.

“We look forward to characterising all foregrounds and then being able to reveal the CMB in unprecedented detail.”

Planck’s first cosmological dataset is expected to be released in 2013.

Provided by European Space Agency

Citation: Planck mission steps closer to the cosmic blueprint (2012, February 13) retrieved 30 April 2024 from <https://phys.org/news/2012-02-planck-mission-closer-cosmic-blueprint.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.