

Scientists deploy lasers, GPS technology to improve snow measurements

February 13 2012



Ethan Gutmann examines a laser instrument for measuring snow. Credit: ©UCAR, Photo by Carlye Calvin

- Equipped with specialized lasers and GPS technology, scientists at the National Center for Atmospheric Research (NCAR) are working with colleagues to solve a critical wintertime weather mystery: how to accurately measure the amount of snow on the ground.

Transportation crews, [water managers](#), and others who make vital safety decisions need [precise measurements](#) of how [snow](#) depth varies across wide areas. But traditional measuring devices such as snow gauges or yardsticks often are inadequate for capturing snow totals that can vary even within a single field or neighborhood.

Now scientists are finding that [prototype devices](#) that use light pulses, satellite signals, and other technologies offer the potential to almost instantly measure large areas of snow. In time, such devices might even provide a global picture of snow depth.

"We've been measuring rain accurately for centuries, but snow is much harder because of the way it's affected by wind and sun and other factors," says NCAR scientist Ethan Gutmann. "It looks like new technology will finally give us the ability to say exactly how much snow is on the ground."

NCAR is conducting the research with several collaborating organizations, including the University of Colorado Boulder and the [National Oceanic and Atmospheric Administration](#) (NOAA). The work is supported by NCAR's sponsor, the National Science Foundation.

Uncertain depths

Emergency managers rely on snowfall measurements when mobilizing snow plows or deciding whether to shut down highways and airports during major storms. They also use snow totals when determining whether a region qualifies for disaster assistance. In mountainous areas, officials need accurate reports of snowpack depth to assess the threat of avalanches or floods, and to anticipate the amount of water available from spring and summer runoff.

More [accurate measurements](#) can also help meteorologists and hydrologists better understand snow physics and hydrological processes.



The three-dimensional features of a snow field above treeline is revealed by laser measurements. The laser, installed by NCAR at a test site in the Rocky Mountains, measures snow at more than 1,000 points across an area almost the size of a football field. Credit: ©UCAR, Image by Ethan Gutmann, NCAR

But traditional approaches to measuring snow can greatly underreport or overreport snow totals, especially in severe conditions. Snow gauges may miss almost a third of the snow in a windy storm, even when they are protected by specialized fencing designed to cut down on the wind's impacts. Snow probes or yardsticks can reveal snow depth within limited areas. But such tools require numerous in-person measurements at different locations, a method that may not keep up with totals during heavy snowfalls.

Weather experts also sometimes monitor the amount of snow that collects on flat, white pieces of wood known as snow boards, but this is a time-intensive approach that requires people to check the boards and clear them off every few hours. The nation's two largest volunteer efforts-The National Weather Service's Cooperative Observer Program and the Community Collaborative Rain, Hail, and Snow Network (CoCoRaHS)-each involve thousands of participants nationwide using snow boards, but their reports are usually filed just once a day.

More recently, ultrasonic devices have been deployed in some of the world's most wintry regions. Much like radar, these devices measure the length of time needed for a pulse of ultrasonic energy to bounce off the

surface of the snow and return to the transmitter. However, the signal can be affected by shifting atmospheric conditions, including temperature, humidity, and winds.

Testing new approaches

The specialized laser instruments under development at NCAR can correct for such problems. Once set up at a location, they can automatically measure snow depth across large areas. Unlike ultrasonic instruments, lasers rely on light pulses that are not affected by atmospheric conditions.



Ethan Gutmann walks past a snow gauge at a research site in Colorado. Credit: ©UCAR, Photo by Carlye Calvin

New tests by Gutmann indicate that a laser instrument installed high above treeline in the Rocky Mountains west of Boulder can measure 10 feet or more of snow with an accuracy as fine as half an inch or better. The instrument, in a little over an hour, measures snow at more than 1,000 points across an area almost the size of a football field to produce a three-dimensional image of the snowpack and its variations in depth.

Gutmann's next step, if he can secure the needed funding, will be to build and test a laser instrument that can measure snow over several square miles. Measuring such a large area would require a new instrument capable of taking over 12,000 measurements per second.

"If we're successful, all of a sudden these types of instruments will reveal a continually updated picture of snow across an entire basin," he says.

One limitation for the lasers, however, is the light pulses cannot penetrate through objects such as trees and buildings. This could require development of networks of low-cost laser installations that would each record snow depths within a confined area. Alternatively, future satellites equipped with such lasers might be capable of mapping the entire world from above.

Gutmann and Kristine Larson, a colleague at the University of Colorado, are also exploring how to use [GPS](#) sensors for snowfall measurements. GPS sensors record both satellite signals that reach them directly and signals that bounce off the ground. When there is snow on the ground, the GPS signal bounces off the snow with a different frequency than when it is bare soil, enabling scientists to determine how high the surface of the snow is above the ground.

Such units could be a cost-efficient way of measuring snow totals because meteorologists could tap into the existing global network of ground-based GPS receivers. However, researchers are seeking to fully understand how both the density of the snow and the roughness of its surface alter GPS signals.

"Our hope is to develop a set of high-tech tools that will enable officials to continually monitor snow depth, even during an intense storm," Larson says. "While we still have our work cut out for us, the technology is very promising."

"I think this technology has great potential to benefit emergency managers and other decision makers, as well as forecasters," Gutmann says.

Provided by National Center for Atmospheric Research

Citation: Scientists deploy lasers, GPS technology to improve snow measurements (2012, February 13) retrieved 30 January 2023 from <https://phys.org/news/2012-02-scientists-deploy-lasers-gps-technology.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.