

Tornado season looms, but forecasting a challenge

February 23 2012, By SETH BORENSTEIN , AP Science Writer



In this Jan. 23, 2012, file photo, residents walk around through the debris of their neighborhood after a tornado ripped through the Trussville, Ala. area. Storm chasers are watching the Southeast as a nasty storm brews with the potential to spin off a batch of tornadoes. Tornado season is just about here and forecasters believe twisters are possible Thursday and Friday. But in reality, it got an early and deadly start in late January when two people were killed by separate twisters in Alabama. The season usually starts in March and then ramps up for the next couple of months, but forecasting tornado seasons is even more imprecise than predicting hurricane seasons. (AP Photo/Butch Dill)

(AP) -- Tornado season is starting, but don't ask meteorologists how bad it will be this spring and summer.

They don't know. They're having a hard enough time getting a fix on the likely path of storms expected in the next 48 hours, from the Ohio Valley to the Southeast coast.

The very nature of tornadoes makes them the wildcard of [weather disasters](#). It's just hard to figure when and where they'll appear.

If funnel clouds do develop by week's end, a real possibility, they won't be the first. In January two people were killed by separate twisters in Alabama. Preliminary reports showed 95 tornadoes struck last month, compared with 16 in January 2011, a particularly stormy year.

The season usually starts in March and then ramps up for the next couple of months, but forecasting a seasonal outlook is even more imprecise than predicting hurricane seasons. Tornadoes are too small and too short-lived. They don't develop like blizzards and hurricanes, which are easier to project.

They pop in and pop out. The storms that give them birth may last only a few hours. Hurricanes and blizzards are lumbering beasts that spend days moving across the satellite maps. When a hurricane approaches, coastlines get days to evacuate. With a tornado, if the [weather service](#) can let people know 20 minutes in advance, it's considered a victory.

"The Joplin (Mo.) tornado (that killed 158 people last May) wasn't violent until just about the time it got to the hospital," said Harold Brooks, a research scientist at the National Oceanic and Atmospheric Administration's National Severe Storms Laboratory, in Norman, Okla. "Even when you're in the field, there are still times when you're surprised by the intensity of the event and how quickly it started."

If a forecast for a hurricane or blizzard is off by a mile, it's still [bad weather](#). But a mile difference means no damage in a tornado, Brooks said: "It's so much finer in time and space on the tornado, it does make it a harder problem."

It takes a piece of debris only a few seconds to fly around an entire

tornado; it takes hours to circle a hurricane. Yet tornadoes, though smaller, can have stronger winds. Since 1950, there have been 58 tornadoes in the United States with winds exceeding 200 mph; six last year alone. Only three hurricanes have made U.S. landfall with winds more than 155 mph.

And forecasters are telling the Southeast and Midwest to get ready again.

"It looks like this week we're moving into a slightly more active dynamic pattern," said warning meteorologist Greg Carbin at the National Weather Service's Storm Prediction Center, also in Norman, Okla.

The percolating Ohio Valley/Southeastern storm is proof of how hard meteorologists have it. On Tuesday evening, Carbin said, "We're kind of expecting it to be a fairly significant event" and the storm center's website had a small red swath for potential severe storms with tornadoes.

By Wednesday afternoon, the storm prediction center massively expanded its Thursday watch area to include 14 states from Florida to Indiana. By Thursday afternoon, it was down to mostly Kentucky and Tennessee with parts of neighboring states.

"A lot of things have to come together at once to have a tornadic storm and the skill at forecasting all those things is near zero," said Howard Bluestein, a professor at the University of Oklahoma. "They are definitely more unpredictable."

All this comes on the heels of one of the worst tornado years in U.S. history. Tornadoes in 2011 started the earliest ever - New Year's Day - killing 550 people, injuring 5,400 and causing \$10 billion in damage over the year, the most in U.S. history. The 2011 season had the most tornadoes in a single day and a single month on record.

But if you ask tornado experts what that means for this year, they'll answer that they just don't know.

Some meteorologists mention La Nina, the flip side of El Nino, as an indicator. It's a cooling of the central Pacific Ocean. Scientists have noticed a correlation between strong La Ninas and active tornado seasons - including last year. But it's not that simple or clear-cut, Columbia University professor Michael Tippett said. The current La Nina is weakening so much it shouldn't be a factor this year, several experts said.

Tippett has a new study that gives some hope, pointing out potential factors - vertical wind shear, updraft and a type of rainfall - that might help for long-range tornado forecasts.

Later this summer, meteorologists will meet in a special conference to try to figure out how to do that type of longer-term tornado prediction. And the National Weather Service is installing new radar for live forecasting, tracking and distinguishing of [tornadoes](#) that could save lives in real-time because forecasters can be more certain in their warnings, said National Weather Service meteorologist Paul Schlatter.

All those elements together mean that maybe by 2020 or so, meteorologists will be able to say watch out this season or relax a bit - but not just yet, said federal researcher Brooks.

More information:

National Weather Service's Storm Prediction Center:

<http://www.spc.noaa.gov/>

National Weather Service's summary of 2011 tornado season:

<http://1.usa.gov/wvq3t8>

©2012 The Associated Press. All rights reserved. This material may not be published, broadcast, rewritten or redistributed.

Citation: Tornado season looms, but forecasting a challenge (2012, February 23) retrieved 26 April 2024 from <https://phys.org/news/2012-02-tornado-season-looms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.