

'Lucy' lived among close cousins: Discovery of foot fossil confirms two human ancestor species co-existed

March 28 2012



(PhysOrg.com) -- A new fossil discovery from Eastern Africa called the Burtele foot indicates *Australopithecus afarensis*, an early relative of modern humans, may not have been the only hominin to walk the plains and woodlands of what is now the Afar region of Ethiopia some 3.4 million years ago.



Researchers openly have questioned whether *Au. afarensis*, the species to which the famous fossil "Lucy" belongs, was the only living hominin during the late Pliocene of Africa. Lucy's bones provided evidence that she and perhaps other early hominins may have walked upright, but whether or not she was the sole hominin species in her particular geologic time scale has been the subject of much debate.

"There was indeed more than one early hominin species during that time," said Yohannes Haile-Selassie, head of Physical Anthropology at the Cleveland Museum of Natural History and lead author of the findings.

Haile-Selassie and a team of anthropologists and geologists report finding the partial skeleton of a foot that belonged to an early human ancestor that was neither *Au. afarensis* nor another hominin called Kenyanthropus platyops, a creature that some paleoanthropologists argue was a second hominin that lived at the same time as *Au. afarensis*.



The journal Nature published the finding today. The National Science



Foundation's (NSF) Division of Behavioral and Cognitive Sciences partially funded the research.

Descriptions of Kenyanthropus platyops led some scientists to dispute whether only one hominin inhabited Africa's upper regions 3.9 to 2.9 million years ago mainly due to the distorted nature of the specimen used as the basis for the original published description of Kenyanthropus platyops, said Haile-Selassie.

"Tim White, (a member of the research team who began uncovering Lucy in 1973), argued that Kenyanthropus platyops is a Kenyan version of *Australopithecus afarensis* and that the subtle differences between the two could be subsumed into an intra-specific variation."

In other words, he argued there was no evidence a second hominin roamed the landscape during the time of *Au. afarensis*. But Haile-Selassie and his team may have found proof of at least two hominin species.

In Woranso-Mille, a relatively new palaeontological site located in the central Afar region of Ethiopia, researchers dug up a 3.4 million year old partial foot skeleton that does not match the contemporaneous *Au*. *afarensis* in form or shape.

Moreover, the skeletal remains infer locomotor adaptations more similar to an earlier, 4.4 million year old hominin, *Ardipithecus ramidus*, that was discovered by a research team led by White in 1992-1993 in the Middle Awash valley in Ethiopia.

"The Burtele foot differs from *Australopithecus afarensis* largely by possessing an opposable great toe," said Haile-Selassie, noting the Burtele toe is more like that of *Ardipithecus ramidus*, inferring similar characteristics for walking, running and jumping.



"This partial pedal skeleton is unique in providing important evidence bearing on the functional morphology and proportions of several early hominin foot elements," the researchers write in their article titled, "A new hominin foot from Ethiopia shows multiple Pliocene bipedal adaptations."

Carolyn Ehardt, program director for Biological Anthropology at NSF, points out that research findings such as this foster appreciation for the complex processes that "shaped the evolutionary history of our species."

"We become increasingly aware of the fact that the evolutionary history of hominins is not unlike that of other groups of organisms in the potential for morphological and behavioral diversity and multiple adaptive pathways characterizing those life-forms during particular evolutionary time periods," she said.



The sifted dirt is carefully picked using small picking pans and each bone fragment is picked and checked for possible fragments of the partial foot. Credit: © The Cleveland Museum of Natural History

The Burtele foot has some skeletal ratios that fall within the human and gorilla distribution, but outside those of chimpanzees. In addition, the



metatarsal of the fourth toe is longer than that of the second toe, a condition seen in some monkeys and Miocene apes.

"Unfortunately this ratio is unknown for both *Ardipithecus* and *Australopithecus*," said Haile-Selassie. "But, the finding could indicate the primitive condition for the human family."

The researchers say identifying and naming the species to which the Burtele foot belongs will have to await recovery of more fossils, cranial and dental elements, for example. However, they are sure it does not belong to the species of Lucy, *Au. afarensis*.

Said Haile-Selassie, "It is probably descended from something like *Ardipithecus ramidus*."

Provided by Cleveland Museum of Natural History

Citation: 'Lucy' lived among close cousins: Discovery of foot fossil confirms two human ancestor species co-existed (2012, March 28) retrieved 27 April 2024 from <u>https://phys.org/news/2012-03-lucy-cousins-discovery-foot-fossil.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.