

# Short-term episodicity of Archaean plate tectonics

March 28 2012

---

Plate tectonics, the dominant process shaping Earth as we know it today, may not have existed throughout Earth's history.

Indeed, the interior of our planet (the mantle) cools progressively, by perhaps 300 degrees Celsius over the past 3.0 billion years.

[Numerical calculations](#) reveal that in Archaean times (4.0-2.5 billion years ago), the mantle was too hot to support stable, long-lived plate tectonics.

Rather, Jean-François Moyen and Jeroen van Hunen suggest that subduction -- a key component of plate tectonics, with cold, rigid plates sinking from the surface down into the [mantle](#) -- was an episodic process, stopping and starting frequently.

Evidence for this episodicity is found in rocks from old geological units such as the Abitibi province of the Canadian Shield, where Moyen and van Hunen describe short, repeated, episodic bursts of subduction related lavas interlayered in non-subduction rocks.

They propose that [plate tectonics](#) started progressively on [Earth](#) by more and more frequent, long-lived, and large-sized subduction events progressively evolving into the stable, large structures observed today.

**More information:** Jean-François Moyen, UMR 6524 CNRS and Université Jean-Monnet, 23 rue du Dr Michelon, 42023 Saint-Etienne,

France., Durham DH1 3LE, UK; and Jeroen van Hunen Durham University. *Geology*. Posted online 26 March 2012; [doi: 10.1130/G32894.1](https://doi.org/10.1130/G32894.1)

Provided by Geological Society of America

Citation: Short-term episodicity of Archaean plate tectonics (2012, March 28) retrieved 23 April 2024 from <https://phys.org/news/2012-03-short-term-episodicity-archaean-plate-tectonics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.