

# Cassini finds Saturn moon Phoebe has planet-like qualities

April 27 2012, By Jia-Rui C. Cook and Dwayne Brown

---



Phoebe's true nature is revealed in startling clarity in this mosaic of two images taken during Cassini's flyby on June 11, 2004. Credit: NASA/JPL/Space Science Institute


(Phys.org) -- Data from NASA's Cassini mission reveal Saturn's moon Phoebe has more planet-like qualities than previously thought.

Scientists had their first close-up look at Phoebe when Cassini began exploring the [Saturn](#) system in 2004. Using data from multiple spacecraft instruments and a computer model of the moon's chemistry, geophysics and geology, scientists found Phoebe was a so-called planetesimal, or remnant planetary building block. The findings appear in the April issue of the Journal Icarus.

"Unlike primitive bodies such as comets, Phoebe appears to have actively evolved for a time before it stalled out," said Julie Castillo-Rogez, a planetary scientist at NASA's Jet Propulsion Laboratory in Pasadena, Calif. "Objects like Phoebe are thought to have condensed very quickly. Hence, they represent building blocks of planets. They give scientists clues about what conditions were like around the time of the birth of planets and their moons."

Cassini images suggest Phoebe originated in the far-off Kuiper Belt, the region of ancient, icy, rocky bodies beyond Neptune's [orbit](#). Data show Phoebe was spherical and hot early in its history, and has denser rock-rich material concentrated near its center. Its average density is about the same as Pluto, another object in the [Kuiper Belt](#). Phoebe likely was captured by Saturn's gravity when it somehow got close to the giant planet.

Saturn is surrounded by a cloud of irregular moons that circle the planet in orbits tilted from Saturn's orbit around the sun, the so-called equatorial plane. Phoebe is the largest of these irregular moons and also has the distinction of orbiting backward in relation to the other moons. Saturn's large moons appear to have formed from gas and dust orbiting in the planet's equatorial plane. These moons currently orbit Saturn in that same plane.



This panel of images shows the nearly spherical shape of Saturn's moon Phoebe, as derived from imaging obtained from NASA's Cassini spacecraft. Each image represents a 90-degree turn. Credit: NASA/JPL-Caltech/SSI/Cornell

"By combining Cassini data with modeling techniques previously applied to other solar system bodies, we've been able to go back in time and clarify why it is so different from the rest of the Saturn system," said Jonathan Lunine, a co-author on the study and a Cassini team member at Cornell University, Ithaca, N.Y.

analyses suggest that Phoebe was born within the first 3 million years of the birth of the solar system, which occurred 4.5 billion years ago. The moon may originally have been porous but appears to have collapsed in on itself as it warmed up. Phoebe developed a density 40 percent higher than the average inner Saturnian moon.

Objects of Phoebe's size have long been thought to form as "potato-shaped" bodies and remained that way over their lifetimes. If such an object formed early enough in the solar system's history, it could have harbored the kinds of radioactive material that would produce substantial heat over a short timescale. This would warm the interior and reshape the moon.

"From the shape seen in Cassini images and modeling the likely cratering history, we were able to see that Phoebe started with a nearly spherical shape, rather than being an irregular shape later smoothed into a sphere by impacts," said co-author Peter Thomas, a Cassini team member at Cornell.

Phoebe likely stayed warm for tens of millions of years before freezing up. The study suggests the heat also would have enabled the moon to host liquid water at one time. This could explain the signature of water-rich material on Phoebe's surface previously detected by Cassini.

The new study also is consistent with the idea that several hundred million years after Phoebe cooled, the moon drifted toward the inner [solar system](#) in a solar-system-wide rearrangement. Phoebe was large enough to survive this turbulence.

More than 60 moons are known to orbit Saturn, varying drastically in shape, size, surface age and origin. Scientists using both ground-based observatories and Cassini's cameras continue to search for others.

The Cassini-Huygens mission is a cooperative project of NASA, the European Space Agency and the Italian Space Agency. JPL manages the mission for the agency's Science Mission Directorate in Washington. The California Institute of Technology in Pasadena manages JPL for NASA.

**More information:** [saturn.jpl.nasa.gov](http://saturn.jpl.nasa.gov)

Provided by JPL/NASA

Citation: Cassini finds Saturn moon Phoebe has planet-like qualities (2012, April 27) retrieved 18 April 2024 from <https://phys.org/news/2012-04-cassini-saturn-moon-phoebe-planet-like.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.