

## **'Cosmic mirages' confirm accelerated cosmic expansion**

April 11 2012



Figure 1 Image of SDSSJ1226-0006, a new gravitationally lensed quasar discovered in this survey. The quasar image in the original image of the Sloan Digital Sky Survey (SDSS), which has been used for the actual survey to identify gravitational lensing, looks only slightly extended, but the Hubble Space Telescope (HST) image clearly exhibits two distinct quasar images (white) as well as a massive galaxy in between the quasar images (orange) that produces gravitational lensing.



(Phys.org) -- An international team of researchers led by Masamune Oguri at Kavli IPMU and Naohisa Inada at Nara National College of Technology conduced an unprecedented survey of gravitationally lensed quasars, and used it to measure the expansion history of the universe. The result provides strong evidence that the expansion of the universe is accelerating. There were several observations that suggested the accelerated cosmic expansion, including distant supernovae for which the 2011 Nobel Prize in Physics was awarded. The team's result confirms the accelerated cosmic expansion using a completely different approach, which strengthens the case for dark energy. This result will be published in *The Astronomical Journal*.

Quasars are very luminous objects powered by <u>accretion</u> of gas into <u>supermassive black holes</u> at the centers of distant galaxies. A quasar is typically located far away. "<u>Gravitational lensing</u>, a phenomenon that a <u>distant object</u> is split into two or more images due to the gravity of a massive foreground object." Toshifumi Futamase, Professor at <u>Tohoku</u> University says, "Prof. Oguri is a leading young researcher in the research field of gravitational lensing. He leads the test of the presence of dark energy using gravitational lensing. The great advantage of this research is much higher statistical significance, which is made possible by a large sample of <u>quasars</u> in the SDSS." The phenomenon of gravitational lensing, often called as "Cosmic Mirage", was first discovered in 1979, and since then more than 100 gravitationally lensed quasars have been reported.

An international team of researchers led by Masamune Oguri, Assistant Professor at the Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU), and Naohisa Inada, Lecturer at Nara National College of Technology, conducted a large survey to search for gravitationally lensed quasars in the massive data sets of the Sloan Digital Sky Survey (SDSS). Over almost 10 years of careful examinations of 100,000 quasars, the team successfully discovered



nearly 50 new gravitationally lensed quasars in total, significantly increased a sample of cosmic mirages (Figure 1).



Figure 2 Illustration of the gravitational lensing measurement of the cosmic expansion speed. The accelerated expansion increases the distance to the quasar, giving rise to higher chance of having a massive galaxy very close to the light path to produce gravitational lensing.

The frequency of gravitational lensing, which can be measured by counting the number of gravitationally lensed quasars within a given quasar catalog, allows one to infer the expansion speed of the universe, because the accelerated expansion increases the distance to each quasar and therefore enhances the chance of gravitational lensing (Figure 2).

The team measured the probability of gravitational lensing among distant quasars to be about 0.05 percent, which was then compared with detailed theoretical calculations to extract information on the expansion hisotry.



The result indicates that the <u>expansion of the universe</u> is indeed accelerating, which suggests that the universe must be filled with mysterious energy component called dark energy.

"The accelerated <u>cosmic expansion</u> is one of the central problems in modern cosmology", Oguri says, "In 2011, the <u>Nobel Prize in Physics</u> was awarded to the discovery of the accelerated expansion of the universe using observations of distant supernovae. A caution is that this method using supernovae is built on several assumptions, and therefore independent checks of the result are important in order to draw any robust conclusion. Our new result using gravitational lensing not only provides additional strong evidence for the accelerated cosmic expansion, but also is useful for accurate measurements of the expansion speed, which is essential for investigating the nature of dark energy."

Careful comparisons with other cosmological observation led to the conclusion that dark energy behaves almost like Einstein's cosmological constant. "Statistical method on gravitationally lensed quasars have been known to be sensitive to the expansion history of the universe," Masashi Chiba, Professor at Tohoku University, says,"therefore actively studied by Japanese researchers in the 1990s. Observations of gravitational lensing at that time already hinted the presence of dark energy, but both due to the small sample size and large uncertainty in the theoretical modeling of lensing rates the result was not widely accepted. This research conducted a enormous survey of gravitationally lensed quasars and adopted much more sophisticated theoretical calculations to build a very convincing case for the accelerated cosmic expansion."

"Studies of dark energy (cosmological constant) were popular in Japan already in the early 1990s, largely because of pioneering work by Prof. Fukugita, Principal Investigator at Kavli IPMU and a co-author of this paper.", says Yasushi Suto, Professor at University of Tokyo, "This survey of gravitationally lensed quasars was initiated and organized by



Profs. Oguri and Inada, who were graduate students when the survey started, within the large international SDSS collaboration, and they led the team to success. This result is important in that it confirms the presence of dark energy independently from the observation of supernovae."

Detailed properties of dark energy are planned to be explored in the SuMIRe project, an international survey project led by Kavli IPMU using the Subaru telescope. Suto says, "This result demonstrates that Japanese theoretical and observational cosmologists will play an essential role in the SuMIRe project." Chiba also says, "This result creates big momentum for the survey of the cosmic <u>dark energy</u> by the SuMIRe project."

**More information:** *The Astronomical Journal*, "The Sloan Digital Sky Survey Quasar Lens Search. VI. Constraints on Dark Energy and the Evolution of Massive Galaxies," Masamune Oguri, Naohisa Inada et al. Preprint: <u>arxiv.org/abs/1203.1088</u>

Provided by Kavli IPMU

Citation: 'Cosmic mirages' confirm accelerated cosmic expansion (2012, April 11) retrieved 17 May 2024 from <u>https://phys.org/news/2012-04-cosmic-mirages-expansion.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.