

# **Researchers find new form of Mars lava flow**

#### April 26 2012



Image: Arizona State University

High-resolution photos of lava flows on Mars reveal coiling spiral patterns that resemble snail or nautilus shells. Such patterns have been found in a few locations on Earth, but never before on Mars. The discovery, made by Arizona State University graduate student Andrew Ryan, is announced in a paper published April 27, 2012, in the scientific journal *Science*.

The new result came out of research into possible interactions of <u>lava</u> <u>flows</u> and floods of water in the Elysium volcanic province of <u>Mars</u>.

"I was interested in Martian outflow channels and was particularly



intrigued by Athabasca Valles and Cerberus Palus, both part of Elysium," says Ryan, who is in his first year as a graduate student in ASU's School of Earth and Space Exploration, part of the College of Liberal Arts and Sciences. Philip Christensen, Regents' Professor of Geological Sciences at ASU, is second author on the paper.

"Athabasca Valles has a very interesting history," Ryan says. "There's an extensive literature on the area, as well as an intriguing combination of seemingly fluvial and volcanic features." Among the features are large slabs or plates that resemble broken floes of pack ice in the Arctic Ocean on Earth. In the past, a few scientists have argued that the plates in Elysium are in fact underlain by water ice.

## Looking below

Assessing those claims that ice was present today beneath the lava plates drove Ryan to study the area. "My initial goal," he says, "was to model the nighttime infrared temperatures of the plates. Then I became fascinated by the terrain lying between the plates and the high-centered polygonal patterns found there." This led him to look closely at every available image of the region.





Image: Arizona State University

"I examined probably 100 HiRISE images of the area," he says, referring to the <u>High Resolution Imaging Science Experiment</u> camera on the Mars Reconnaissance Orbiter. In addition, he pored over daytime and nighttime infrared and visual images from the Thermal Emission Imaging System (THEMIS) camera on Mars Odyssey orbiter. (Christensen is THEMIS' principal investigator.) Images taken by the Context Camera (CTX) on <u>Mars Reconnaissance Orbiter</u>, the High Resolution Stereo Camera (HRSC) on Mars Express, and the Mars Orbiter Camera (MOC) on Mars Global Surveyor were all studied as well.



## **Picture this**

"One evening," Ryan recounts, "I was making a second pass over the HiRISE images when I first noticed puzzling spiral patterns in an image near the southern margin of Cerberus Palus. I actually nearly overlooked that particular frame, thinking that it might not be too useful, being so far away from main study area farther north."

He notes, "The coils become noticeable in the full-resolution HiRISE image only when you really zoom in. They also tend to blend in with the rest of the light-gray terrain — that is, until you stretch the contrast a bit.

"I don't find it surprising that these were overlooked in the past. I nearly missed them too."

## **Curling lava**

On Earth, lava coils can be found on the Big Island of Hawaii, mainly on the surface of ropey pahoehoe lava flows. They have also been seen in submarine lava flows near the Galapagos Rift on the Pacific Ocean floor.

As Ryan explains, "The coils form on flows where there's a shear stress — where flows move past each other at different speeds or in different directions. Pieces of rubbery and plastic lava crust can either be peeled away and physically coiled up — or wrinkles in the lava's thin crust can be twisted around."

Similarly, he notes that scientists have documented the formation of rotated pieces of oceanic crust at mid-ocean ridge spreading centers. "Since the surface of active lava lakes, such as those on Hawaii, can have crustal activity like spreading centers do, it's conceivable that lava coils



may form there in a similar way, but at a smaller scale."

The size of Martian lava coils came as a surprise. "On Mars the largest lava coil is 30 meters across — that's 100 feet. That's bigger than any known lava coils on Earth," he says. Ryan and Christensen's work has inventoried nearly 200 lava coils in the Cerberus Palus region alone.

Looking ahead, Ryan says, "Lava coils may be present in other Martian volcanic provinces or in outflow channels mantled by volcanic features. I expect that we'll find quite a few more in Elysium as the <u>HiRISE</u> image coverage grows over time."

Provided by Arizona State University

Citation: Researchers find new form of Mars lava flow (2012, April 26) retrieved 26 April 2024 from <u>https://phys.org/news/2012-04-mars-lava.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.