

## Mathematics: First-ever image of a flat torus in 3D

April 26 2012



Image showing the isometric embedding of a square flat torus in 3D space, seen from the outside (above) and from the inside (below). Different oscillation waves, called corrugations, can be distinguished. Together, the corrugations form an object that resembles a fractal and has a rough appearance. © Borrelli, Jabrane, Lazarus, Thibert

Just as a terrestrial globe cannot be flattened without distorting the distances, it seemed impossible to visualize abstract mathematical objects called flat tori in ordinary three-dimensional space. However, a French team of mathematicians and computer scientists has succeeded in constructing and visually representing an image of a flat torus in three-dimensional space. This is a smooth fractal, halfway between fractals and ordinary surfaces. The results are published in *PNAS*.



In the 1950s, Nicolaas Kuiper and the <u>Nobel laureate</u> John Nash demonstrated the existence of a representation of an abstract mathematical object called flat torus, without being able to visualize it. Since then, constructing a representation of this surface has remained a challenge that has finally been met by scientists in Lyon and Grenoble.

On the basis of the Convex Integration Theory developed by Mikhail Gromov in the 1970s, the researchers used the corrugation technique (oscillations). This reputedly abstract <u>mathematical method</u> helps to determine atypical solutions to partial differential equations. This enabled the scientists to obtain images of a flat torus in 3D for the first time. Halfway between fractals and ordinary surfaces, these images show a smooth <u>fractal</u>.

These findings open up new avenues in applied mathematics, especially in the visualization of the differential equations encountered in physics and biology. The astounding properties of smooth fractals could also play a central role in the analysis of the geometry of shapes.

**More information:** Flat tori in three dimensional space and convex integration - Vincent Borrelli, Saïd Jabrane, Francis Lazarus and Boris Thibert, *Proceedings of the National Academy of Sciences (PNAS)*, April 2012.

To find out more about this research: <u>www.gipsa-lab.fr/~francis.lazarus/Hevea/Presse/</u>

Provided by CNRS

Citation: Mathematics: First-ever image of a flat torus in 3D (2012, April 26) retrieved 26 April 2024 from <u>https://phys.org/news/2012-04-mathematics-first-ever-image-flat-torus.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.