

## **Researchers develop method to fingerprint air pollution**

## April 20 2012

A team of researchers from the University of Pittsburgh and the Electric Power Research Institute (EPRI) collected emissions samples from several power plant stacks in the United States and developed a unique method for detecting the isotopic signatures of nitrogen oxide (NOx) emissions under different configurations. These isotopic signatures will be instrumental in helping to identify emission sources of air pollution across the nation.

NOx <u>emissions</u> are formed during the combustion of fossil fuels in, for instance, internal combustion engines or power plants. NOx emissions mix with organic gases in the atmosphere to form <u>ozone</u> and particulate matter, the main components of <u>smog</u>. These emissions eventually settle onto surfaces, and the deposited material, primarily nitrate, carries a measurable isotopic signature. However, until now, scientists were unable to fully interpret these signatures because they lacked the "fingerprints" of various NOx emission sources.

Pitt and EPRI researchers developed a method of extracting NOx emission samples from one of these sources—the stacks of coal-fired power plants—and measuring their isotopic composition. Sampling took place at facilities with and without advanced NOx-reducing technologies. Researchers discovered that emissions from power plants employing the advanced NOx controls had different proportions of the 15N atom in the NOx they emitted than the plants without the advanced technologies or NOx emissions from other sources. With this information, scientists will be able to analyze deposition samples and



better determine the sources contributing to the deposited NOx products.

"We've been mapping the isotopes of nitrogen oxide deposition products across the nation," said principal investigator Emily Elliott, assistant professor in the Pitt Kenneth P. Dietrich School of Arts and Sciences Department of Geology and Planetary Sciences. "These 'isoscapes' can only be interpreted with fingerprint data like the isotopic signatures collected in this study."

Pitt's researchers modified existing EPA methods to collect NOx from power plant stacks. Once collected, Elliott's lab utilized bacteria to convert nitrate into a gaseous form for isotopic analysis. Prior to the coupling of these techniques, previous analytical approaches were both time- and labor-intensive and precluded widespread characterization of environmental nitrate isotopes. The "Regional Stable Isotope Laboratory for Earth and Environmental Research" study directed by Elliott is now using these methods to examine isoscapes of nitrogen in air, water, and across ecosystems.

"Stable isotopes have been used successfully in science for a variety of purposes," said Stephanie Shaw, senior project manager at EPRI. "The characterization of NOx isotope geochemistry will allow researchers to determine how different sources and different emission-control technologies actually influence air quality."

These results, combined with additional information from other NOx sources, will allow scientists to look at rain samples and determine how much nitrogen comes from <u>power plants</u> stacks as opposed to how much comes from such other sources as motor vehicles, lightning, or soil. J. David Felix, a project team member and doctoral candidate at Pitt, is now working to identify the isotopic composition of other reactive nitrogen emission sources, including those produced in animal feed lots, fertilizer applications, and emissions produced by vehicles. The team is



conducting a pilot study with Pitt's School of Public Health to examine the isotopic composition of <u>nitrogen oxides</u> and determine their sources within the city of Pittsburgh, where exposures relevant to human health may be occurring.

Since incorporating low-NOx combustors and emissions control technologies like SCR, power plant emissions have decreased by more than 40 percent since 2005 and nearly 70 percent since 1990. Power plant NOx emissions are expected to continue to decline in future years.

"Based on these results, the overall isotopic composition of power plant NOx emissions is expected to change and thus change the isotopic composition of nitrogen in environmental samples," Elliott stated. "It was important for us to understand how the implementation of emission control technologies affects the isotopic nature of the NOx being emitted in order to evaluate its fate in the atmosphere."

The study was published in the March 2012 issue of *Environmental Science and Technology*.

Provided by University of Pittsburgh

Citation: Researchers develop method to fingerprint air pollution (2012, April 20) retrieved 1 May 2024 from <u>https://phys.org/news/2012-04-method-fingerprint-air-pollution.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.