


Study finds warm ocean currents cause majority of ice loss from Antarctica

April 25 2012



This image shows the circulation of ocean currents around the western Antarctic ice shelves. The shelves are indicated by the rainbow color; red is thicker (greater than 550 meters), while blue is thinner (less than 200 meters). Credit: NASA/Goddard CGI Lab

Reporting this week in the journal *Nature*, an international team of scientists led by British Antarctic Survey (BAS) has established that warm ocean currents are the dominant cause of recent ice loss from Antarctica. New techniques have been used to differentiate, for the first time, between the two known causes of melting ice shelves - warm ocean currents attacking the underside, and warm air melting from above. This finding brings scientists a step closer to providing reliable projections of future sea-level rise.

Researchers used 4.5 million measurements made by a laser instrument mounted on NASA's ICESat satellite to map the changing thickness of

almost all the [floating ice](#) shelves around Antarctica, revealing the pattern of ice-shelf melt across the continent. Of the 54 ice shelves mapped, 20 are being melted by [warm ocean](#) currents, most of which are in [West Antarctica](#).

In every case, the inland glaciers that flow down to the coast and feed into these thinning ice shelves have accelerated, draining more ice into the sea and contributing to sea level rise.

Lead author Dr Hamish Pritchard from [British Antarctic Survey](#), which is part of the UK's Natural Environment Research Council (NERC), said:

"In most places in Antarctica, we can't explain the ice-shelf thinning through melting of snow at the surface, so it has to be driven by warm ocean currents melting them from below.

"We've looked all around the Antarctic coast and we see a clear pattern: in all the cases where ice shelves are being melted by the ocean, the inland glaciers are speeding up. It's this glacier acceleration that's responsible for most of the increase in ice loss from the continent and this is contributing to sea-level rise."

"What's really interesting is just how sensitive these glaciers seem to be. Some ice shelves are thinning by a few metres a year and, in response, the glaciers drain billions of tons of ice into the sea. This supports the idea that [ice shelves](#) are important in slowing down the glaciers that feed them, controlling the loss of ice from the Antarctic [ice sheet](#). It means that we can lose an awful lot of ice to the sea without ever having summers warm enough to make the snow on top of the glaciers melt - the oceans can do all the work from below.

"But this does raise the question of why this is happening now. We think

that it's linked to changes in wind patterns. Studies have shown that Antarctic winds have changed because of changes in climate, and that this has affected the strength and direction of [ocean currents](#). As a result warm water is funnelled beneath the floating ice. These studies and our new results therefore suggest that Antarctica's [glaciers](#) are responding rapidly to a changing climate."

A different picture is seen on the eastern Antarctic Peninsula (the long stretch of land pointing towards South America). Here, the ice-shelf thinning found by this study can be explained by warm summer winds directly melting the snow on the ice-shelf surfaces. Both patterns, of widespread ocean-driven melting and this summer melting on the Antarctic Peninsula, can therefore be attributed to Antarctica's changing wind patterns.

This research is part of international efforts to improve understanding of the interactions between ice and climate in order to improve the reliability of [sea-level rise](#) projections. Professor David Vaughan is the leader of ice2sea - a major EU-funded FP7 programme. He said,

"This study shows very clearly why the Antarctic ice sheet is currently losing ice, which is a major advance. But the real significance is that it also shows the key to predicting how the ice sheet will change in the future is in understanding the oceans. Perhaps we should not only be looking to the skies above Antarctica, but also into the surrounding oceans."

The study was carried out by an international team from British Antarctic Survey, Utrecht University, Scripps Institution of Oceanography and Earth & Space Research in Corvallis, Oregon. NASA's ICESat – Ice, Cloud and Land Elevation Satellite – measurements were collected during the period 2003 – 2008 to detect changes in ice-shelf thickness through time.

More information: Antarctic ice sheet loss driven by basal melting of ice shelves: H.D Pritchard, S.R.M Ligtenberg, H.A Fricker, D.G Vaughan, M.R van den Broeke, L. Padman is published this week in the journal *Nature*.

Provided by British Antarctic Survey

Citation: Study finds warm ocean currents cause majority of ice loss from Antarctica (2012, April 25) retrieved 27 April 2024 from <https://phys.org/news/2012-04-ocean-currents-majority-ice-loss.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--