

# Researchers report potential for a 'moderate' New England 'red tide' in 2012

April 5 2012

---

New England is expected to experience a "moderate" regional "red tide" this spring and summer, report NOAA-funded scientists working in the Gulf of Maine to study the toxic algae that causes the bloom. The algae in the water pose no direct threat to human beings, however the toxins they produce can accumulate in filter-feeding organisms such as mussels and clams— which can cause paralytic shellfish poisoning (PSP) in humans who consume them.

Under a newly developed rating system, a moderate bloom could cause the closure of shellfish beds along an estimated 126 – 250 miles of coastline.

The 2012 outlook is based on the quantities of the algae *Alexandrium fundyense* in its dormant – or cyst – state detected in [Gulf of Maine](#) sediments last fall. These data are combined with computer simulations that model a complex range of meteorological and oceanographic conditions -- winds, sunlight, rainfall, tides, and currents -- that impact the size of the bloom.

"Our goal over the last ten years has been to develop a system to help the shellfish industry and environmental managers better plan for the annual bloom," said Woods Hole Oceanographic Institution (WHOI) biologist Don Anderson, who has worked with WHOI colleague Dennis McGillicuddy and North Carolina State University (NCSU) Prof. Ruoying He to develop the computer model to help predict the intensity and location of blooms.

Scientists are unable to make a precise forecast of where and when the regional bloom will make landfall because bloom transport depends on episodic weather events and currents that cannot be predicted months in advance. Rather, the scientists use the computer model to produce a range of bloom scenarios – or an "ensemble forecast" – tracking variables like wind direction and water characteristics based on previous years' conditions. This is similar to the system used to forecast hurricanes.

A number of factors could impact the forecast. For instance, changing characteristics of water in the Gulf of Maine can have a direct effect on the growing conditions for *Alexandrium*.

"The surveys of cyst abundance gives us an indication of the potential extent of the bloom, but whether or not that potential is realized depends on the growing conditions," said McGillicuddy. "In 2010 we forecast a large bloom but we got it wrong. That spring, an unusual mass of warm, fresh water that was low in nutrients changed the growing conditions."

Wind direction imposes another uncertainty to the forecast. For example, strong northeast winds in the spring and early summer drive the bloom inland toward coastal shellfish beds. In contrast, when southwesterlies dominate, the algae tend to stay offshore.

"Each year, we add another set of environmental conditions to our archive of model runs. In the future, a winter that is warmer and drier than normal can be represented by 2012, but right now, we have no similar year in that archive," said Anderson.

In order to protect public health, shellfish beds are closed when toxicities rise above a quarantine level, often during the peak harvesting season. Due to effective monitoring by state agencies, there have been no illnesses from legally harvested shellfish in recent years, despite some

severe blooms during that time period. There have been, however, several severe poisonings of individuals who ignored closure signs.

The economic impacts of PSP toxicity are significant in the region. Direct and indirect costs of the extensive *Alexandrium* bloom in 2005 were estimated at nearly \$50 million for Massachusetts and \$23 million for Maine.

The 2012 designation of a "moderate" bloom now has a specific, quantifiable meaning, thanks to a complementary research effort by Anderson and his colleagues to develop forecast terminology to describe a bloom's potential impact. As part of that work, Judy Kleindinst, a member of Anderson's team, analyzed records of harvesting closures due to PSP extending back 35 years, and identified three categories of bloom severity.

The categories -- "limited", "moderate", and "extensive" -- are equivalent to closures over 0 - 125, 126 - 250, and 251 - 375 miles of coastline. A moderate outbreak might stretch from Maine to northern Massachusetts, although it could be shifted down the coast, covering the same length of coastline, but over a different area.

When combined, the forecast terminology, the annual cyst surveys and the continual improvements and additions to the computer model have developed into a useful management tool.

"[Red tide](#) is a chronic problem affecting commercial and recreational harvesting interests throughout the Gulf of Maine," said Chris Nash, shellfish program manager for the New Hampshire Department of Environmental Services. "State agencies are responsible for monitoring toxicity levels in shellfish harvest areas and implementing harvest closures to prevent illness outbreaks. These regional scale, seasonal outlooks help state managers to plan and use limited monitoring

resources effectively. Ultimately our goals are to protect public health and give consumers confidence in the quality of the seafood products they purchase from markets and restaurants, and these forecasts are useful in realizing those goals."

"NOAA is developing a HAB Operational Forecasting System (HAB OFS) in order to provide advanced warnings like this one to help state agencies monitor harmful algal blooms and minimize public health risks," said David Kennedy, assistant NOAA administrator for the National Ocean Service. "The Gulf of Maine is one of several regions for which HAB forecasts are being developed with the intent to operationalize them within NOAA utilizing multiple assets such as those provided by the National Weather Service."

Project researchers regularly share their field observations and models with more than 80 coastal resource and fisheries managers in six states as well as federal agencies like NOAA, the EPA, and the FDA. Real-time forecasts are updated on a weekly basis and additional information will be provided on the "Current Status" page of the Northeast PSP website, particularly as new information becomes available from coastal ocean observing systems such as the Northeastern Regional Association of Coastal Ocean Observing Systems (NERACOOS), EcoMon, and AZMP. The National Weather Service is also providing extended hydrological and meteorological outlooks to accompany the bloom forecasts.

Provided by Woods Hole Oceanographic Institution

Citation: Researchers report potential for a 'moderate' New England 'red tide' in 2012 (2012, April 5) retrieved 11 July 2024 from <https://phys.org/news/2012-04-potential-moderate-england-red-tide.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.