

The presence of water changes the structure of an antibiotic

April 20 2012



Part of the gramicidin molecule © INRI

EPFL chemists have shown how the three-dimensional shape of an antibiotic changes when it is in an aqueous environment. This could lead to a better understanding of how drugs interact with biological molecules.

Like a key inserted into a lock, the <u>molecules</u> in drugs bind with and act upon biomolecules. The more precisely we know these molecules' threedimensional structure, the better we will understand how their active components work. Biomolecules often exist in aqueous environments; this is the case in the human body. But water isn't just a neutral environment; it can also interact with the molecules and change their structure. Scientists from EPFL's Laboratory of Molecular Physical



Chemistry have shown how the 3D structure of gramicidin – a natural antibiotic – changes depending on the number of water molecules surrounding it. This research could lead to progress in designing new drugs that could be used against diseases such as epilepsy, cancer and Alzheimer's disease.

To determine the structure of a molecule, chemists measure its spectrum, or the wavelengths of light that it absorbs when illuminated by a laser. This spectral signature allows them to predict the 3D structure of the molecule. Ideally, each spectrum corresponds to a particular structure. But at room temperature, large molecules like gramicidin – made up of 176 atoms – are victims of a parasitic quantum effect known as inhomogeneous thermal broadening. This interference, associated with the thermal agitation of the atoms in the molecule, makes it impossible to get a precise enough measurement to be able to identify a unique molecular structure.

Quantum cold

The chemists had to reduce the effects of these perturbations. To do this, they trapped the molecules in a specially designed device and then quickly cooled the molecules to about 10 degrees Kelvin. At this extremely cold temperature – just ten degrees above absolute zero – the parasitic quantum effects diminish and measurements can be made. Finally, they added water to the antibiotic, molecule by molecule, and observed that its structure changed as a function of the quantity of water molecules added. They were able to make measurements by adding up to fifty molecules of water at a time.

This research, led by senior scientist Oleg Boyarkine, his colleague Natalia Nagornova and professor Thomas Rizzo, will have a considerable impact not only in fundamental research, but also in the pharmaceutical arena. "Understanding how <u>drug</u> molecules change shape



when they are dissolved in water is a crucial point. This discovery will make it possible to design new, more effective drugs using computers," the chemist explains. The challenge was to conduct the experiments as accurately and precisely as possible. "Even though precise knowledge of the structure of a large molecule is only really possible through theoretical calculations, thanks to spectroscopy and ultra-low temperatures, we can now validate the theory."

Treating Alzheimer's disease and epilepsy

Other molecules will be examined using this technique. Drugs developed to combat Alzheimer's disease and epilepsy attempt to short-circuit certain chemical reactions in the brain. The drugs bind with proteins in the brain at specific points, and block disease progression. The creation of this chemical bond is largely dependent on the structure of both the active components in the drug and the target proteins. The goal of future research will be to determine the specific binding points between a drug and a protein, as well as their 3D structures in the presence of water. Understanding these structures will thus open up avenues for developing new drugs.

More information: Science, 19 april 2012.

Provided by Ecole Polytechnique Federale de Lausanne

Citation: The presence of water changes the structure of an antibiotic (2012, April 20) retrieved 2 May 2024 from <u>https://phys.org/news/2012-04-presence-antibiotic.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.