

Size matters - even for molecules

April 11 2012



Two electrons that are emitted from a large molecule by a single photon may originate from far apart within that molecule. In a recent study on hydrocarbon molecules consisting of one to five fused benzene rings (each ring consisting of six carbon atoms), SRC researchers Tim Hartman and Ralf Wehlitz have found that the relative probability for ejecting two electrons scales linearly with the length of the molecule.

This indicates that the two electrons can originate at the two opposite ends of the molecule, which, in this case, is up to 1.4 nm or about 10 times the diameter of a carbon atom apart. This observation also implies that the production of stable doubly charged parent ions relative to singly



charged parent ions is rather large for large molecules leading to the emission of many slow electrons.

When a slow electron attaches to another molecule, such as a large biomolecule, it makes it prone to <u>radiation damage</u>, so that such "marked" molecule can easily fragment.

The publication of this paper by SRC Researchers Tim Hartman and Ralf Wehlitz can be found at <u>T. Hartman et al., *PRL* 108, 023001 (2012)</u>

Provided by University of Wisconsin-Madison

Citation: Size matters - even for molecules (2012, April 11) retrieved 10 May 2024 from https://phys.org/news/2012-04-size-molecules.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.