

When stellar metallicity sparks planet formation

April 10 2012, By Ray Sanders



Artist's concept showing a young Sun-like star surrounded by a planet-forming disk of gas and dust. Credit: NASA/JPL-Caltech/T. Pyle

New research predicts the criteria needed for Earth-like planets to form around a star that have one-tenth the metallicity of our Sun. If researchers find small, rocky planets orbiting stars with lower metallicity, it may challenge the presently accepted "core accretion" model of planetary formation.

In new research, scientists have attempted to determine the precise conditions necessary for planets to form in a <u>star system</u>. Jarrett Johnson and Hui Li of Los Alamos National Laboratory assert that observations increasingly suggest that <u>planet formation</u> takes place in star systems



with higher metallicities.

Astronomers use the term "<u>metallicity</u>" in reference to elements heavier than hydrogen and helium, such as oxygen, silicon, and iron. In the "core <u>accretion</u>" model of <u>planetary formation</u>, a rocky core gradually forms when <u>dust grains</u> that make up the disk of material that surrounds a young star bang into each other to create small rocks known as "planetesimals". Citing this model, Johnson and Li stress that heavier elements are necessary to form the dust grains and planetesimals which build planetary cores.

Additionally, evidence suggests that the circumstellar disks of dust that surround young <u>stars</u> don't survive as long when the stars have lower metallicities. The most likely reason for this shorter lifespan is that the light from the star causes clouds of dust to evaporate.

The Planet Epoch

Our cosmic history has several defining "epochs", one of which is the point at which star systems began to form planets. Heavy elements such as carbon, silicon and oxygen first needed to be created from supernovae and the stellar cores of the first generations of stars before the first planets could form.

"Our calculation is an estimate of the minimum amount of heavy elements that must be present in circumstellar disks before planets can form," says Johnson. "Because these heavy elements must be produced by the first stars in the universe, the first planets could only form around later generations of stars."

Understanding how the first planets formed provides crucial information about the early universe. Additionally, a better understanding of early planetary formation impacts many facets of astronomy, including the



search for life elsewhere.



HST image of a Dust Disk Around Star HD141569. Credit: NASA, M. 1Clampin (STScI), H. Ford (JHU), G. Illingworth (UCO/Lick), J. Krist (STScI), D. Ardila (JHU), D. Golimowski (JHU), the ACS Science Team and ESA

According to the team, a successful theory of planet formation should make predictions about the properties of the earliest planets and their host stars. Such a theory could be tested by studying very old planetary systems in our galaxy. The enrichment of gas with metals from supernovae is thought to not only affect planetary formation, but the formation of low-mass stars like our Sun as well.

"A planet as massive and dense as the <u>Earth</u> could only form once stars and supernovae had enriched the gas with an abundance of heavy elements that is at least 10 percent that in the Sun," adds Johnson. "This suggests that many generations of stars had to form and evolve before habitable planets could form."



One important consideration for planetary formation is the dispersal rate of the circumstellar disk of gas and dust around a host star. Two of the more prominent mechanisms for dispersing a planetary disk are giant planet formation and photoevaporation by the host star. Photoevaporation appears to be the more dominant process which dictates the lifetime of a planetary disk around a star. Observations show that low-metallicity disks have shorter lifetimes, which is bolstered by data showing higher-metallicity disks are better "shielded" from evaporation by a host star's radiation.

Johnson and Li further state that disks with higher metallicity tend to form a greater number of high mass giant planets.



This image shows the critical metallicity for Earth-like planet formation, expressed as iron abundance relative to that of the Sun, as a function of distance (r) from the host star. If systems with planets are discovered in the "forbidden zone", it may pose a challenge to the "core accretion" model of planetary formation. Credit: Johnson & Li

The Lifetime of Dust

In order to obtain estimates of the critical metallicity necessary for



planet formation, Johnson and Li compared the lifetime of the disk and the length of time required for dust grains in the disk to settle. Basically, for a star system to form planets, the time required for dust grains to settle cannot exceed the lifetime of the planetary disk.

The team explains that the dust-to-gas ratio that occurs when the timescales are equal gives an estimate of the critical metallicity, the point at which their model suggests planets can form. Since the settling time for dust grains depends on the density and temperature of the disk, which are related to the distance from the host star, the critical metallicity is also a function of distance from the host star.

"Our calculation is really fairly simple compared to many others, as we have focused only on what we believe are the key processes that set the timescale required for planetesimal formation at low metallicity," Johnson says. "These are the growth of dust grains into planetesimals and the destruction of the disk by the high energy radiation from the host star. While the calculation is simple, it does show that current models of planet formation can in principle explain how the lowest metallicity planets form."

The team notes several assumptions made in their comparisons with the data. The first assumption is that surface metallicity of the host star is the same as that of the protostellar disk from which it and its planets formed. Secondly, the team assumes circular planetary orbits. When orbits are highly eccentric, comparing the data to the theoretical predictions is more difficult. Lastly, the team assumes planets have not migrated inward toward their star from their initial place of birth in the disk.





Mercury transit of the Sun on November 8, 2006. A sunspot located just below the equator at the left-hand side is much bigger than Mercury, which looks like a small black dot in the lower middle of the solar disk.

The team found that the formation of planetesimals can only take place once a minimum metallicity is reached in a protostellar disk. Since the earliest stars that formed in the universe (Population III stars) do not have the required metallicity to host planets, it is believed that the supernova explosions from such stars helped enrich subsequent (Population II) stars, some of which may still be in existence and could host planets.

The Earliest Planets

Based on their equations, the team finds that some of the earliest planets may have formed at a distance of 0.03 AU from their parent star (for comparison, Mercury orbits at just under 0.4 AU). Given the high temperatures at such compact orbits (estimated at roughly 1600 K or 1,300 C), planet formation is likely to have resulted in planets too hot to



host life as we know it.

"Interestingly, our results also suggest that the first Earth-like planets may have formed in the habitable zones of stars slightly more massive than the Sun, Johnson adds. "Because more massive stars burn out faster, it is possible that any life that evolved on these planets may have already perished with the death of its host star, which may have lived only 4 billion years compared to the 10 billion year lifetime expected for the <u>Sun</u>."

Johnson and Li also note that the formation of Earth-like <u>planets</u> is not itself a sufficient prerequisite for life to take hold, stating that early galaxies contained numerous supernovae and black holes - both strong sources of radiation that would threaten life. Given the hostile conditions in the early universe, it is expected that conditions suitable for life were only present after early galaxy formation.

"However, with the wealth of new exoplanets being discovered and characterized, our theory of the minimum metallicity for planet formation may yet be challenged," Johnson concludes. "It will be exciting to see how [our model] holds up."

Johnson and Li's research is scheduled to appear in the *Astrophysical Journal*.

Source: Astrobio.net

Citation: When stellar metallicity sparks planet formation (2012, April 10) retrieved 2 May 2024 from <u>https://phys.org/news/2012-04-stellar-metallicity-planet-formation.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.