


New type of biosensor is fast, super-sensitive

May 29 2012, By Bob Beale



Schematic: Magnets are used to assemble nanoparticles, coated with antibodies to enrofloxacin, between two electrodes: when antibodies leave the nanoparticles the electrical resistance falls.

(Phys.org) -- A whole new class of biosensor that can detect exceptionally small traces of contaminants in liquids in just 40 minutes has been developed by a UNSW-led team of researchers.

Known as a biochemiresistor, it meets a long-standing challenge to create a sensor that is not only super-sensitive to the presence of [chemical compounds](#) but responds quickly. It has countless potential uses for detecting drugs, toxins and [pesticides](#) for biomedical or environmental analysis.

In a paper published in the prestigious chemistry journal *Angewandte Chemie* the researchers describe how they successfully tested the new sensor by detecting tiny traces in milk of the veterinary antibiotic enrofloxacin. The journal has singled out the study for attention as a “Very Important Paper”. Only 5% of papers published by the journal are so designated.

“Enrofloxacin is an antibiotic used in the agricultural industry that can be transferred to the food chain,” notes co-author Scientia Professor Justin Gooding, of the UNSW School of Chemistry and the Australian Centre for Nanomedicine.

“Our biochemiresistor was able to detect enrofloxacin in neat milk in 40 minutes, at level as low as one nanogram in a litre of milk. To put that number in perspective, a nanogram is a billionth of a gram and is the mass of a single cell.

“While that is impressive enough, the sensor is a general concept that can be widely applied across many different fields.”

A biosensor is a portable analytical device that uses biological molecules to detect selectively just one compound within a mix of many others. Small biosensors are already in daily use testing the safety of drinking water, for checking diabetic blood-sugar levels and for pregnancy tests

The biochemiresistor uses gold-coated magnetic nanoparticles modified with antibodies that are selective for the chemical constituent – or analyte - of interest. The nanoparticles are dispersed into the sample for analysis and if the analyte is present some of the antibodies detach from the nanoparticles.

Using a magnet, the nanoparticles are then assembled into a film between two electrodes and the electrical resistance is measured. The

more analyte is present, the more antibodies leave the nanoparticles and the lower the resistance in the nanoparticle film.

“This new type of biosensor is rapid in response because the magnetic nanoparticle biosensors go and get the analyte rather than the usual approach of waiting for the analyte to find the sensing surface,” says Gooding.

“The biochemiresistor is also more sensitive than the usual [biosensor](#) because, as the nanoparticles are dispersed throughout the sample, the entire sample is analysed, not just a small portion of the solution.”

The study’s lead author is Leo M.H. Lai.

More information: [onlinelibrary.wiley.com/doi/10 ...
e.201202350/abstract](https://onlinelibrary.wiley.com/doi/10.1002/anie.201202350/abstract)

Provided by University of New South Wales

Citation: New type of biosensor is fast, super-sensitive (2012, May 29) retrieved 24 April 2024 from <https://phys.org/news/2012-05-biosensor-fast-super-sensitive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.