

Black hole caught red-handed in a stellar homicide

May 7 2012



(Phys.org) -- Astronomers have gathered the most direct evidence yet of a supermassive black hole shredding a star that wandered too close.

Supermassive black holes, weighing millions to billions times more than the Sun, lurk in the centers of most galaxies. These hefty monsters lay quietly until an unsuspecting victim, such as a star, wanders close enough to get ripped apart by their powerful gravitational clutches.

[Astronomers](#) have spotted these stellar [homicides](#) before, but this is the first time they can identify the victim. Using a slew of ground- and space-based telescopes, a team of astronomers led by Suvi Gezari of The Johns Hopkins University in Baltimore, Md., has identified the victim as a star rich in [helium gas](#). The star resides in a galaxy 2.7 billion light-years away.

Her team's results will appear May 2 in the online edition of the journal *Nature*.

"When the star is ripped apart by the [gravitational forces](#) of the black hole, some part of the star's remains falls into the black hole, while the rest is ejected at high speeds. We are seeing the glow from the stellar gas falling into the black hole over time. We're also witnessing the spectral signature of the ejected gas, which we find to be mostly helium. It is like we are gathering evidence from a crime scene. Because there is very little hydrogen and mostly helium in the gas we detect from the carnage, we know that the slaughtered star had to have been the helium-rich core of a stripped star," Gezari explained.

This observation yields insights about the [harsh environment](#) around [black holes](#) and the types of [stars](#) swirling around them.

This is not the first time the unlucky star had a brush with the behemoth black hole. Gezari and her team think the star's hydrogen-filled envelope surrounding its core was lifted off a long time ago by the same black hole. In their scenario, the star may have been near the end of its life. After consuming most of its [hydrogen fuel](#), it had probably ballooned in size, becoming a red giant. The astronomers think the bloated star was looping around the black hole in a highly elliptical orbit, similar to a comet's elongated orbit around the Sun. On one of its close approaches, the star was stripped of its puffed-up atmosphere by the black hole's powerful gravity. Only its core remained intact. The stellar remnant

continued its journey around the black hole, until it ventured even closer to the behemoth monster and faced its ultimate demise.

Astronomers have predicted that stripped stars circle the central black hole of our Milky Way galaxy, Gezari pointed out. These close encounters, however, are rare, occurring roughly every 100,000 years. To find this one event, Gezari's team monitored hundreds of thousands of galaxies in ultraviolet light with the NASA's Galaxy Evolution Explorer (GALEX), a space-based observatory, and in visible light with the Pan-STARRS1 telescope on the summit of Haleakala in Hawaii. Pan-STARRS, short for Panoramic Survey Telescope and Rapid Response System, scans the entire night sky for all kinds of transient phenomena, including supernovae.

The team was looking for a bright flare in ultraviolet light from the nucleus of a galaxy with a previously dormant black hole. They found one in June 2010, which was spotted with both telescopes. Both telescopes continued to monitor the flare as it reached peak brightness a month later and then slowly began to fade over the next 12 months. The brightening event was similar to that of a supernova, but the rise to the peak was much slower, taking nearly one and a half months.

"The longer the event lasted, the more excited we got, since we realized that this is either a very unusual supernova or an entirely different type of event, such as a star being ripped apart by a black hole," said team member Armin Rest of the Space Telescope Science Institute in Baltimore, Md.

By measuring the increase in brightness, the astronomers calculated the black hole's mass to be several million suns, which is comparable to the size of our Milky Way's black hole.

Spectroscopic observations with the MMT (Multiple Mirror Telescope)

Observatory on Mount Hopkins in Arizona showed that the black hole was swallowing lots of helium. Spectroscopy divides light into its rainbow colors, which yields an object's characteristics, such as its temperature and gaseous makeup.

"The glowing helium was a tracer for an extraordinarily hot accretion event," Gezari said. "So that set off an alarm for us. And, the fact that no hydrogen was found set off a big alarm that this was not typical gas. You can't find gas like that lying around near the center of a galaxy. It's processed gas that has to have come from a stellar core. There's nothing about this event that could be easily explained by any other phenomenon."

The observed speed of the gas also linked the material to a black hole's gravitational pull. MMT measurements revealed that the gas was moving at more than 20 million miles an hour (over 32 million kilometers an hour). However, measurements of the speed of gas in the interstellar medium reveal velocities of only about 224,000 miles an hour (360,000 kilometers an hour).

"The place we also see these kinds of velocities are in supernova explosions," Rest said. "But the fact that it is still shining in ultraviolet light is incompatible with any supernova we know."

To completely rule out the possibility of an active nucleus flaring up in the galaxy, the team used NASA's Chandra X-ray Observatory to study the hot gas. Chandra showed that the characteristics of the gas didn't match those from an active galactic nucleus.

"This is the first time where we have so many pieces of evidence, and now we can put them all together to weigh the perpetrator (the black hole) and determine the identity of the unlucky star that fell victim to it," Gezari said. "These observations also give us clues to what evidence to

look for in the future to find this type of event."

Provided by JPL/NASA

Citation: Black hole caught red-handed in a stellar homicide (2012, May 7) retrieved 10 April 2024 from <https://phys.org/news/2012-05-black-hole-caught-red-handed-stellar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.