


Environmental group measures methane seeps in the Arctic

May 21 2012, by Bob Yirka



The effect on lake ice formation of subcap and superficial seeps. Image (c)
Nature Geoscience (2012) doi:10.1038/ngeo1480

(Phys.org) -- A team of researchers, led by Katey Walter Anthony, of the University of Alaska, has been studying and mapping so-called seeps, holes in lake ice near the edges of glaciers where methane is bubbling up from below into the atmosphere, and suggest that a feedback loop may be in the works. As the team writes in their paper describing their observations, which has been published in the journal *Nature Geoscience*, seeping methane that results from Arctic warming, might itself also be contributing to warming; a feedback loop that could accelerate global warming.

Methane is considered by most Earth scientists to be number two on the

list of [greenhouse gasses](#) that contribute to global warming, after CO₂, and because of that more research is being done with the hopes of finding where all of it is coming from. It generally forms when organic matter decays or is digested and thus known sources include animal exhaust, including from humans, melting of permafrost, waste from [landfills](#) and decaying matter beneath the floor of the world's oceans. Now, based on this new research it appears another significant source is seeps from lakes formed by glaciers.

Scientists believe that when glaciers formed they trapped organic matter beneath them. Later as lakes formed at their edges, methane began to be produced as the [organic matter](#) rotted. That methane remained trapped however, until recently, by the cap of ice that remained on the surface of such lakes. Now, in the past decade, with the rise in Arctic temperatures, seeps have developed allowing the methane to slowly escape into the atmosphere. The team has mapped, using both aerial and ground based surveys, 150,000 such seeps in Alaska and Greenland alone.

Scientists monitoring methane in the atmosphere have found that levels of the gas rose predictably in the latter half of the last century, then for unknown reasons, leveled off for about five years, and then resumed rising again. Because of this, researchers are concerned, as it's thought that the Arctic, which is experiencing warming at a faster pace than anywhere else on the planet, could be entering a phase where because of feedback loops such as these seen with [methane](#) (and with carbon emissions from melting permafrost) we're on a course that could see global warming occur on a much faster pace than has been previously anticipated. Others of course suggest that because we still don't really understand how the whole system works, making any sort of prediction at this point would be premature.

More information: Geologic methane seeps along boundaries of Arctic permafrost thaw and melting glaciers, *Nature Geoscience* (2012)

[doi:10.1038/ngeo1480](https://doi.org/10.1038/ngeo1480)

Abstract

Methane, a potent greenhouse gas, accumulates in subsurface hydrocarbon reservoirs, such as coal beds and natural gas deposits. In the Arctic, permafrost and glaciers form a ‘cryosphere cap’ that traps gas leaking from these reservoirs, restricting flow to the atmosphere. With a carbon store of over 1,200 Pg, the Arctic geologic methane reservoir is large when compared with the global atmospheric methane pool of around 5 Pg. As such, the Earth’s climate is sensitive to the escape of even a small fraction of this methane. Here, we document the release of ^{14}C -depleted methane to the atmosphere from abundant gas seeps concentrated along boundaries of permafrost thaw and receding glaciers in Alaska and Greenland, using aerial and ground surface survey data and in situ measurements of methane isotopes and flux. We mapped over 150,000 seeps, which we identified as bubble-induced open holes in lake ice. These seeps were characterized by anomalously high methane fluxes, and in Alaska by ancient radiocarbon ages and stable isotope values that matched those of coal bed and thermogenic methane accumulations. Younger seeps in Greenland were associated with zones of ice-sheet retreat since the Little Ice Age. Our findings imply that in a warming climate, disintegration of permafrost, glaciers and parts of the polar ice sheets could facilitate the transient expulsion of ^{14}C -depleted methane trapped by the cryosphere cap.

© 2012 Phys.Org

Citation: Environmental group measures methane seeps in the Arctic (2012, May 21) retrieved 15 May 2024 from <https://phys.org/news/2012-05-environmental-group-methane-seeps-arctic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.