


# Forest diversity from Canada to the subtropics influenced by family proximity

May 17 2012

---



Data from over 3 million trees in the eastern half of the US were aggregated into two-degree-latitude-by-longitude cells in order to study regional patterns of conspecific negative density dependence, a process where the mortality of a species rises in coincidence with its increasing abundance. Credit: Indiana University

How species diversity is maintained is a fundamental question in biology. In a new study, a team of Indiana University biologists has shown for the first time that diversity is influenced on a spatial scale of unparalleled scope, in part, by how well tree seedlings survive under their own parents.

Scientists have long considered conspecific negative density dependence (CNDD), a process where the mortality of a [species](#) rises in coincidence with its increasing abundance, to be a key mechanism maintaining [diversity](#) at the local scale. In new research to be published Friday in the journal *Science*, the IU researchers show that this mechanism is driving diversity from the boreal forests to sub-tropical forests.

The report, "Conspecific negative density dependence and forest diversity," is authored by Daniel Johnson, a doctoral student in the IU Bloomington College of Arts and Sciences' Department of Biology. Co-authors are Wesley T. Beaulieu, also a doctoral student in the Department of Biology, and biology professors James D. Bever and Keith Clay, Johnson's major advisor.

Their work analyzed data on [forest composition](#) from over 200,000 plots containing more than 1.3 million trees and from paired plots containing over 1.7 million seedlings of 151 different tree species. The plots were located from the Canadian border south to Florida and from the Atlantic coast to the 100th meridian and covered over 1.5 million square miles. The U.S. Forest Service spends about \$62 million each year to gather the publicly available forest inventory data used in the IU study.

"We are now able to provide robust evidence that CNDD is pervasive in forest communities from boreal to sub-tropical regions and that it can significantly affect the relative abundance and richness of species with and between forests," Johnson said. "And we now see that the ability to which one tree species can sustain itself in the same area has profound impacts on the diversity of species at a spatial scale that has not been attainable previously. This is the first time it's been shown to be happening not just at a local spatial scale but over the entire eastern US."

The concept of CNDD is based on the well-known Janzen-Connell hypothesis, which proposes that the close proximity of adults reduces

seedling survival of that species through increased attack by host-specific pests and pathogens.

Studies of CNDD in the past have mostly focused on forest communities at single sites or of a single species, with the most recent work showing that in [tree species](#), composition and abundance can be influenced by CNDD at the scale of individual trees.

"Local interactions have previously been considered to affect [species diversity](#) at a local scale, but our findings indicate that local interactions feed back to species richness and abundance over much larger geographical scale, spanning most of eastern North America," Johnson said.

Evidence that local interactions underlie regional species richness is in contrast to the current understanding that patterns of forest diversity are primarily driven by temperature, precipitation and other physical aspects of the environment. This discovery has implications for how [forest](#) modeling is conducted and conservation and management decisions are made.

Provided by Indiana University

Citation: Forest diversity from Canada to the sub-tropics influenced by family proximity (2012, May 17) retrieved 17 July 2024 from <https://phys.org/news/2012-05-forest-diversity-canada-sub-tropics-family.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.