

Genetic safety in numbers, platypus study finds

May 18 2012



The more genetic diversity in a platypus population, the more protected they are against disease.

(Phys.org) -- Platypuses on the Australian mainland and in Tasmania are fighting fit but those on small islands are at high risk of being wiped out from disease, according to a University of Sydney study.

The finding has important implications, not only for the management of the [platypus](#) but for other populations with limited [genetic variation](#), including the iconic koala.

"We have identified platypus populations which are at high risk of disease epidemics due to low [genetic diversity](#)," said Mette Lillie, a PhD researcher at the University of Sydney's Faculty of [Veterinary Science](#).

Lillie is the lead author of the study on platypus populations published this month in the [Journal of Heredity Advance Access](#). Lillie's supervisor Kathy Belov, Associate Professor in Animal Genetics is the study's senior author.

The platypus has a wide distribution in Australia, including island populations on Tasmania, King Island and [Kangaroo Island](#).

King Island is located north-west of the main [island of Tasmania](#) and Kangaroo Island is south-west of Adelaide.

"Our study measured how immunologically fit platypus populations are based on how much variation there is in their immune genes. Variation is very important for populations because the more variation there is, the more pathogens the populations can resist," Lillie said.

The study found platypus colonies on the mainland and in Tasmania show high immune gene diversity. The smaller island platypus populations were a different story. The Kangaroo Island population had very low diversity and the King Island population had no immune gene diversity at all.

The lack of diversity in the Kangaroo Island population can be explained by the fact that only a small number of platypuses were introduced there during the 1930s and 40s, which would have limited genetic diversity.

The King Island population has been isolated from Tasmania and the mainland since the [last ice age](#), 14,000 years ago, and genetic diversity would have been lost in this small population due to inbreeding and random genetic drift an important evolutionary force that influences genetic diversity from one generation to the next. Genetic drift is especially significant in small isolated populations where it tends to decay genetic diversity over time.

The team has previously shown that low immune gene diversity allowed a contagious cancer to decimate Tasmanian devils. Identification of a shallow gene pool in Kangaroo and King Island platypuses raises concerns about their long term survival.

These populations will need to be carefully managed and monitored for signs of disease outbreaks, as an introduced disease could reach epidemic proportions.

This research has important implications for conservation management as it reinforces the idea that healthy populations need high levels of immune gene diversity. Large populations of immunological clones are a recipe for disaster, and time bombs for disease epidemics.

Provided by University of Sydney

Citation: Genetic safety in numbers, platypus study finds (2012, May 18) retrieved 11 May 2024 from <https://phys.org/news/2012-05-genetic-safety-platypus.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.