

Scientists discover first ever record of insect pollination from 100 million years ago

May 14 2012



This is a synchrotron tomography image of the specimen of *Gymnospollisthrips minor* studied at the ESRF. The pollen grains are highlighted in yellow. Credit: ESRF

Amber from Cretaceous deposits (110-105 my) in Northern Spain has revealed the first ever record of insect pollination. Scientists have discovered in two pieces of amber several specimens of tiny insects covered with pollen grains, revealing the first record of pollen transport and social behavior in this group of animals. The results are published in the *Proceedings of the National Academy of Sciences (PNAS)* dated 14-18 May 2012.

The international team of scientists comprises: Enrique Peñalver and Eduardo Barrón from the Instituto Geológico y Minero de España in Madrid; Xavier Delclòs from the University of Barcelona; Andre and Patricia Nel from the Muséum national d'histoire naturelle in Paris; Conrad Labandeira from the Smithsonian Institution, Washington DC; and Carmen Soriano and Paul Tafforeau from the European Synchrotron Radiation Facility (ESRF) in Grenoble, France. The amber samples were from the collection of the Museo de Ciencias Naturales de Álava (Spain).

Today, more than 80% of plant species rely on insects to transport [pollen](#) from male to female flower parts. Pollination is best known in flowering plants but also exists in so-called gymnosperms, seed-producing plants like conifers. Although the most popular group of pollinator insects are bees and butterflies, a myriad of lesser-known species of flies, beetles or thrips have co-evolved with plants, transporting pollen and in return for this effort being rewarded with food.



This is a photo taken under a microscope of Gymnosperm pollen, attached to the abdomen and wing of a *Gymnopollisthrips* fossilized insect inclusion in the Alava amber. Credit: Enrique Peñalver/IGME

During the last 20 years, amber from the Lower Cretaceous (110-105 my) found in the Basque country in Northern Spain has revealed many new plant and animal species, mainly insects. Here, the amber featured inclusions of thysanopterans, so-called thrips, a group of minute insects of less than 2 mm in length that feed on pollen and other plant tissues. They are efficient pollinators for several species of flowering plants.

Two amber pieces revealed six fossilised [specimens](#) of female thrips with hundreds of pollen grains attached to their bodies. These insects exhibit highly specialized hairs with a ringed structure to increase their ability to collect pollen grains, very similar to the ones of well known pollinators like domestic bees. The scientists describe these six specimens in a new genus (*Gymnopollisthrips*) comprising two new species, *G. minor* and *G. major*.

The most representative specimen was also studied with synchrotron X-ray tomography at the ESRF to reveal in three dimensions and at very high resolution the [pollen grain](#) distribution over the insect's body.

The pollen grains are very small and exhibit the adherent features needed so that insects can transport them. The scientists conclude that this pollen is from a kind of cycad or ginkgo tree, a kind of living fossil of which only a few species are known to science. Ginkgos trees are either male or female, and male trees produce small pollen cones whereas female trees bear ovules at the end of stalks which develop into seeds after pollination.

For which evolutionary reason did these tiny insects, 100 million years ago, collect and transport Gingko pollen? Their ringed hairs cannot have grown due to an evolutionary selection benefitting the trees. The benefit for the thrips can only be explained by the possibility to feed their larvae with pollen. This suggests that this species formed colonies with larvae living in the ovules of some kind of gingko for shelter and protection, and female insects transporting pollen from the male Gingko cones to the female ovules to feed the larvae and at the same time pollinate the trees.

Only amber can preserve behavioural features like pollination in such rich detail over millions of years. 100 million years ago, flowering plants started to diversify enormously, eventually replacing conifers as the dominant species. "This is the oldest direct evidence for pollination, and the only one from the age of the dinosaurs. The co-evolution of flowering plants and insects, thanks to pollination, is a great evolutionary success story. It began about 100 million years ago, when this piece of amber fossil was produced by resin dropping from a tree, which today is the oldest fossil record of pollinating [insects](#). Thrips might indeed turn out to be one of the first pollinator groups in geological history, long before evolution turned some of them into flower pollinators", concludes Carmen Soriano, who led the investigation of the amber pieces with X-ray tomography at the ESRF.

Provided by European Synchrotron Radiation Facility

Citation: Scientists discover first ever record of insect pollination from 100 million years ago (2012, May 14) retrieved 19 April 2024 from <https://phys.org/news/2012-05-scientists-insect-pollination-million-years.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.