

# Social status promotes faster wound healing in wild baboons

May 21 2012

---



(c) Wikipedia

Turns out it's not bad being top dog, or in this case, top baboon.

A new study by University of Notre Dame biologist Beth Archie and colleagues from Princeton University and Duke University finds that high-ranking male baboons recover more quickly from injuries and are less likely to become ill than other males.

Archie, Jeanne Altman of Princeton and Susan Alberts of Duke examined [health records](#) from the Amboseli Baboon Research Project in Kenya. They found that high rank is associated with faster wound healing. The finding is somewhat surprising, given that top-ranked males also experience high stress, which should suppress immune responses.

They also found that social status is a better predictor of [wound healing](#) than age.

"In humans and animals, it has always been a big debate whether the stress of being on top is better or worse than the stress of being on the bottom," Archie, lead researcher on the study, said. "Our results suggest that, while animals in both positions experience stress, several factors that go along with high rank might serve to protect males from the negative [effects of stress](#)."

"The power of this study is in identifying the [biological mechanisms](#) that may confer health benefits to high-ranking members of society," George Gilchrist, program director in the National Science Foundation (NSF)'s Division of Biology, which funded the research. "We know that humans have such benefits, but it took meticulous long-term research on baboon society to tease out the specific mechanisms. The question remains of causation: is one a society leader because of stronger immune function or vice versa?"

The researchers examined 27 years of data on naturally occurring illness and injuries in wild male [baboons](#), which is a notably large data set. Although research in health and disease in animals in laboratory settings has been quite extensive, this study is one of most comprehensive ever conducted on animals in a natural setting.

The research team investigated how differences in age, physical condition, stress, reproductive effort and testosterone levels contribute to status-related differences in immune functions. Previous research found that high testosterone levels and intense reproductive efforts can suppress immune function and are highest among high-ranking males.

However, Archie and her colleagues found that high-ranking males were less likely to become ill and recovered faster from injuries and illnesses

than low-ranking males. The authors suggest that chronic stress, old age and poor physical condition associated with low rank may suppress [immune function](#) in low-ranking males.

"The complex interplay among social context, physiology and immune system-mediated health costs and benefits illustrates the power of interdisciplinary research," Carolyn Ehardt, NSF program director for biological anthropology, which co-funded the research. "This research begins to tease apart the trade-offs in both high and low status in primates, including ourselves, which may lead to understanding the effects of [social status](#) on death and disease — not inconsequential for society as a whole."

**More information:** "Social status predicts wound healing in wild baboons," by Elizabeth A. Archie, Jeanne Altmann, and Susan C. Alberts, *PNAS* (2012).

Provided by University of Notre Dame

Citation: Social status promotes faster wound healing in wild baboons (2012, May 21) retrieved 23 May 2024 from <https://phys.org/news/2012-05-social-status-faster-wound-wild.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.