

Dinosaurs were lighter than previously thought, new study shows

June 5 2012



Scientists have developed a new technique to accurately measure the weight and size of dinosaurs and discovered they are not as heavy as previously thought.

University of Manchester [biologists](#) used lasers to measure the minimum amount of skin required to wrap around the skeletons of modern-day [mammals](#), including reindeer, [polar bears](#), giraffes and [elephants](#).

They discovered that the [animals](#) had almost exactly 21% more [body mass](#) than the minimum skeletal 'skin and bone' wrap volume, and applied this to a giant Brachiosaur [skeleton](#) in Berlin's Museum für Naturkunde.


Previous estimates of this Brachiosaur's weight have varied, with estimates as high as 80 tonnes, but the Manchester team's calculations – published in the journal *Biology Letters* – reduced that figure to just 23 tonnes. The team says the new technique will apply to all dinosaur weight measurements.



Brachiosaurus scan

Lead author Dr Bill Sellers said: "One of the most important things palaeobiologists need to know about fossilised animals is how much they weighed. This is surprisingly difficult, so we have been testing a new approach. We laser scanned various large mammal skeletons, including polar bear, giraffe and elephant, and calculated the minimum wrapping volume of the main skeletal sections.

"We showed that the actual volume is reliably 21% more than this value, so we then [laser](#) scanned the Berlin Brachiosaur, *Giraffatitan brancai*, calculating the skin and bone wrapping volume and added 21%. We found that the giant herbivore weighed 23 tonnes, supporting the view that these animals were much lighter than traditionally thought.



Dr Sellers, based in Manchester's Faculty of Life Sciences, explained

that body mass was a critical parameter used to constrain biomechanical and physiological traits of organisms.

He said: "Volumetric methods are becoming more common as techniques for estimating the body masses of fossil vertebrates but they are often accused of excessive subjective input when estimating the thickness of missing soft tissue.

"Here, we demonstrate an alternative approach where a minimum convex hull is derived mathematically from the point cloud generated by laser-scanning mounted skeletons. This has the advantage of requiring minimal user intervention and is therefore more objective and far quicker.

"We tested this method on 14 large-bodied mammalian skeletons and demonstrated that it consistently underestimated body mass by 21%. We suggest that this is a robust method of estimating body mass where a mounted skeletal reconstruction is available and demonstrate its usage to predict the body mass of one of the largest, relatively complete sauropod dinosaurs, *Giraffatitan brancai*, as 23,200 kg.

"The value we got for *Giraffatitan* is at the low range of previous estimates; although it is still huge, some of the enormous estimates of the past – 80 tonnes in 1962 – are exaggerated. Our method provides a much more accurate measure and shows [dinosaurs](#), while still huge, are not as big as previously thought."

More information: *Images: WI Sellers & PL Manning*

Provided by University of Manchester

Citation: Dinosaurs were lighter than previously thought, new study shows (2012, June 5)
retrieved 20 April 2024 from
<https://phys.org/news/2012-06-dinosaurs-lighter-previously-thought.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.