


Internal cellular sensors make *Salmonella* dangerous: study

June 15 2012, By Bill Hathaway



Salmonella bacteria (in green) are engulfed by immune system cell called a macrophage. The direct threat from the immune cell activates genes that trigger the organism to become virulent and dangerous to the host.

(Phys.org) -- *Salmonella* becomes dangerously virulent only when molecular sensors within the organism sense changes in the environment, a team of researchers from the Yale School of Medicine and the Yale Microbial Diversity Institute report in the June 14 issue of the journal *Nature*.

Other [dangerous pathogens](#) probably possess analogous sensors to activate [virulence genes](#) and cause maladies such as [food poisoning](#) and tuberculosis, the authors suggest. This mechanism may present a novel

target for drugs that can disarm bacteria's ability to cause disease, said Eduardo A. Groisman, professor of microbial pathogenesis at the Yale School of Medicine and an investigator with the Howard Hughes Medical Institute.

“There will never be a world without [Salmonella](#) because it exists in many, many animal reservoirs,” Groisman said. “So you can try to avoid getting Salmonella or learn how to fight it. Hundreds of thousands of people die from Salmonella poisoning each year, and tens of millions of people are infected. It is a major public health issue.”

Groisman and Eun-Jin Lee of Yale investigated whether signals from outside the bacteria triggered one of its virulence genes. The researchers meticulously tracked the molecular chain reaction that occurs after Salmonella becomes engulfed by macrophages, immune system cells that respond to bacterial invaders. Salmonella then can reproduce rapidly, often overwhelming defenses of the host.

The Nature paper discovers that it is not the direct threat from the environment itself that triggers the virulence gene within the bacterium. Instead, changes in the level of acidity in Salmonella's surroundings trigger an increase in levels of ATP, the energy currency of all cells. It is the change in ATP levels within the organism that activates virulence factors and enables Salmonella to survive within its host.

This internal sensor acts in concert with several other factors necessary before these bacteria can become virulent, Groisman said. Understanding all these processes should help scientists develop defenses against infectious diseases, he adds.

Provided by Yale University

Citation: Internal cellular sensors make Salmonella dangerous: study (2012, June 15) retrieved 30 April 2024 from

<https://phys.org/news/2012-06-internal-cellular-sensors-salmonella-dangerous.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.