

Research duo discover why non-Newtonian fluids harden on impact

July 12 2012, by Bob Yirka

(Phys.org) -- Researchers and laymen alike know that some non-Newtonian fluids tend to harden quickly upon impact. Quicksand is a good example. If a person walks quickly, they won't sink. But why this happens has been somewhat of a mystery, though many in the science community have suspected it had something to do with energy being transferred to the walls of the container. New research finds that's not the case at all. Instead, after extensive testing and experimentation Scott Waitukaitis and Heinrich Jaeger of the University of Chicago have found, as they describe in their paper published in the journal *Nature*, that it's because the particles suspended in the liquid temporarily mash together, creating a near solid.

Non-Newtonian liquids are those that don't behave as Isaac Newton originally theorized, e.g. quicksand, ketchup, custard, blood, or in the case of the material studied by the research team, a water and cornstarch solution called [oobleck](#).

To find out what goes on with oobleck when impacted from above, Waitukaitis set up several different sized tanks of the goo and filmed what happens when a flat bottomed rod strikes its surface from above at varying speeds. He also pointed an x-ray gun at the mix to see what happens underneath the surface and used a laser pointed across the surface to measure how its surface changed when impacted along with a force sensor on the bottom of the tank directly beneath the impact zone.

In studying the results, the two found that the size of the tank didn't

matter as the oobleck hardened the same amount regardless. Thus, theories about energy absorption by the walls accounting for the unusual liquid properties are wrong. Instead they found that the tiny particles that are normally suspended in the liquid are suddenly jammed together when impacted from above, creating a cone like shape inside the liquid that is dense enough to be described as a temporary solid; as it just as quickly dissolves back to its original state. The authors describe the action as akin to snow that is suddenly pushed into a denser state by a plow.

The study does answer the major question of why a non-Newtonian liquid hardens, but still a mystery is what happens with the suspended particles when in their hardened state. Do they touch each other or just come close and why do they move apart again after the initial impact? Also, are there differences in hardening as particles get smaller and smaller? Future researchers will no doubt be looking into answering such questions as non-Newtonian liquids might become more useful if their properties become better understood. They might serve as a lighter material in bullet-proof vests for example, or provide a better cushion for people involved in a car crash.

More information: Impact-activated solidification of dense suspensions via dynamic jamming fronts, *Nature* 487, 205–209 (12 July 2012). [doi:10.1038/nature11187](https://doi.org/10.1038/nature11187)

Abstract

Although liquids typically flow around intruding objects, a counterintuitive phenomenon occurs in dense suspensions of micrometre-sized particles: they become liquid-like when perturbed lightly, but harden when driven strongly. Rheological experiments have investigated how such thickening arises under shear, and linked it to hydrodynamic interactions or...

See also: [Press release](#)

© 2012 Phys.org

Citation: Research duo discover why non-Newtonian fluids harden on impact (2012, July 12)
retrieved 30 April 2024 from

<https://phys.org/news/2012-07-duo-non-newtonian-fluids-harden-impact.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
