


# Researchers measure the electrical charge of nano particles

July 30 2012



This is a cross-section through two chip-sized glass plates in which a nano particle is trapped in an energy hole (or “potential well” to use the scientific term). The colored fields show the different charges in the electrostatic field. The red zone signifies a very low charge, while the blue edges have a strong charge. Credit: University of Zurich

Nano particles are a millionth of a millimeter in size, making them invisible to the human eye. Unless, that is, they are under the microscope of Prof. Madhavi Krishnan, a biophysicist at the University of Zurich. Prof. Krishnan has developed a new method that measures not only the size of the particles but also their electrostatic charge. Up until now it

has not been possible to determine the charge of the particles directly.

In order to observe the individual [particles](#) in a solution, Prof. Madhavi Krishnan and her co-workers «entice» each particle into an "electrostatic trap". It works like this: between two glass plates the size of a chip, the researchers create thousands of round energy holes. The trick is that these holes have just a weak [electrostatic charge](#). The scientists then add a drop of the solution to the plates, whereupon each particle falls into an energy hole and remains trapped there. But the particles do not remain motionless in their trap. Instead, molecules in the solution collide with them continuously, causing the particles to move in a circular motion. "We measure these movements, and are then able to determine the charge of each individual particle," explains Prof. Madhavi Krishnan.

Put simply, particles with just a small charge make large circular movements in their traps, while those with a high charge move in small circles. This phenomenon can be compared to that of a light-weight ball which, when thrown, travels further than a heavy one. The US physicist Robert A. Millikan used a similar method 100 years ago in his oil drop experiment to determine the velocity of electrically charged oil drops. In 1923, he received the Nobel Prize in physics in recognition of his achievements. "But he examined the drops in a vacuum," Prof. Krishnan explains. "We on the other hand are examining nano particles in a solution which itself influences the properties of the particles."

## **Electrostatic charge of 'nano drugs packages'**

For all solutions manufactured industrially, the electrical charge of the nano particles contained therein is also of primary interest, because it is the electrical charge that allows a fluid solution to remain stable and not to develop a lumpy consistency. "With our new method, we get a picture of the entire suspension along with all of the particles contained in it," emphasizes Prof. Madhavi Krishnan. A suspension is a fluid in which

miniscule particles or drops are finely distributed, for example in milk, blood, various paints, cosmetics, vaccines and numerous pharmaceuticals. "The charge of the particles plays a major role in this," the Zurich-based scientist tells us.

One example is the manufacture of medicines that have to be administered in precise doses over a longer period using drug-delivery systems. In this context, [nano particles](#) act as «packages» that transport the drugs to where they need to take effect. Very often, it is their [electrical charge](#) that allows them to pass through tissue and cell membranes in the body unobstructed and so to take effect. «That's why it is so important to be able to measure their charge. So far most of the results obtained have been imprecise», the researcher tells us.

"The new method allows us to even measure in real-time a change in the charge of a single entity," adds Prof. Madhavi Krishnan. "This is particularly exciting for basic research and has never before been possible." This is because changes in charge play a role in all bodily reactions, whether in proteins, large molecules such as the DNA double helix, where genetic make-up is encoded, or cell organelles. "We're examining how material works in the field of millionths of a millimeter."

**More information:** Mojarad, N, and Krishnan, M., Measuring the size and charge of single nanoscale objects in solution using an electrostatic fluidic trap. *Nature Nanotechnology* (2012) [doi:10.1038/nnano.2012.99](https://doi.org/10.1038/nnano.2012.99)

Provided by University of Zurich

Citation: Researchers measure the electrical charge of nano particles (2012, July 30) retrieved 1 May 2024 from <https://phys.org/news/2012-07-electrical-nano-particles.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.