

Jets at CMS and the determination of their energy scale

July 11 2012, By Henning Kirschenmann




Figure 1: Sketch of pp-collision and resulting collimated spray of particles, a jet.

Jets are the experimental signatures of quarks and gluons produced in high-energy processes such as head-on proton-proton collisions. As quarks and gluons have a net colour charge and cannot exist freely due to colour-confinement, they are not directly observed in Nature. Instead, they come together to form colour-neutral hadrons, a process called hadronisation that leads to a collimated spray of hadrons called a jet.

As these jets of [particles](#) propagate through the CMS detector, they leave signals in components such as the tracker and the electromagnetic and hadronic calorimeters. These signals are combined using jet algorithms to form a reconstructed jet. However, the energy of the reconstructed jets does not correspond to the true particle-level energy, which is independent of detector response. The jet energy corrections relate these two values; the detailed understanding of the jet energy scale is of crucial importance for many physics analyses, and it is often an important component of their systematic uncertainty.

Within CMS, three different methods to reconstruct jets are supported with jet energy corrections: a calorimeter-based approach; the “Jet-Plus-Track” approach, which improves the measurement of [calorimeter jets](#) by exploiting the associated tracks; and the “Particle Flow” approach, which attempts to reconstruct individually each particle in the event, prior to the jet clustering, based on information from all relevant sub-detectors. The significantly improved resulting jet energy resolutions, especially at low transverse momentum (p_T), indicate that the “Particle Flow” approach offers advantages with respect to the other types. Indeed, it is used by most physics analyses within CMS.

CMS follows a factorised approach to jet energy corrections. At first, the additional energy in the event induced by overlaid low-energy proton-[proton collisions](#) (pile-up) is subtracted. For one high-energy head-on proton-proton collision, there are in 2012 – on average – more than 20 of such additional collisions taking place at the same time.

Secondly, to compensate for the non-linear response of the calorimeters (as a function of p_T) and variations of the response in pseudo-rapidity (η), corrections are determined from simulations. These corrections are cross-checked using data-driven methods: Currently, there are analyses determining the relative scale as a function of η from di-jet events and the absolute scale in the central detector region ($|\eta|$

Citation: Jets at CMS and the determination of their energy scale (2012, July 11) retrieved 25 April 2024 from <https://phys.org/news/2012-07-jets-cms-energy-scale.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
