

Polar bear evolution tracked climate change, new DNA study suggests

July 23 2012



Credit: Susanne Miller, U.S. Fish and Wildlife Service. Public domain image.


A whole-genome analysis suggests that polar bear numbers waxed and waned with climate change, and that the animals may have interbred with brown bears since becoming a distinct species millions of years ago.

An analysis of newly sequenced polar bear genomes is providing important clues about the species' evolution, suggesting that climate change and genetic exchange with brown bears helped create the polar bear as we know it today. The international study, led by the Penn State University and the University at Buffalo, found evidence that the size of the polar bear population fluctuated with key [climatic events](#) over the past million years, growing during periods of cooling and shrinking in warmer times.

The research also suggests that while polar bears evolved into a [distinct species](#) as many as 4-5 million years ago, the animals may have interbred with brown bears until much more recently. These intimate relations may be tied to changes in the Earth's climate, with the retreat of glaciers bringing the two species into greater contact as their ranges overlapped, said Charlotte Lindqvist, the study's senior author and an assistant professor of biology at UB.

"Maybe we're seeing a hint that in really warm times, polar bears changed their life-style and came into contact, and indeed interbred, with brown bears," said Stephan Schuster, co-lead author, a professor of biochemistry and molecular biology at Penn State, and a research scientist at Nanyang Technological University in Singapore.

The findings will be published online in the early edition of the [Proceedings of the National Academy of Sciences](#) on 23 July 2012. The study is the most extensive analysis to date of polar bear DNA, scientists say. The research team, representing 13 institutions in North America, Europe, Asia, and Canada, as well as Mexico's Laboratorio Nacional de Genomica para la Biodiversidad (Langebio), sequenced and analyzed the nuclear genomes of 28 different bears, with many [DNA samples](#) provided by the U.S. Geological Survey and the Norwegian Polar Institute.



As shown in this image, the nuclear genomes of bears (black outline), suggest that polar bears and brown bears diverged from one another 4 to 5 million years ago, and that occasional exchange of genes between the two species (shaded gray areas) followed. Results from maternally inherited mitochondrial DNA (dotted line) indicates extinction (marked with an "X") and replacement of polar bear mitochondrial DNA around 160,000 years ago due to interbreeding between the two species. Credit: Penn State University

"We generated a first-rate set of data, including deep sequence coverage for the entire genomes of a polar bear, three brown bears, and a black bear, plus lower coverage of 23 additional polar bears, including a 120 thousand year old individual; very few vertebrate species have such comprehensive genomic resources available," Schuster said. Using this vast amount of data, the scientists discovered that polar bears are actually an older species than previously thought -- indeed, far more ancient than suggested by a recent study that placed the species' age at 600,000 years old. That analysis looked only at small segments of DNA.

"We showed, based on a consideration of the entire DNA sequence, that earlier inferences were entirely misleading," said study co-lead author Webb Miller, a Penn State professor of biology and computer science and engineering. "Rather than polar bears splitting from brown bears a few hundred thousand years ago, we estimate that the split occurred 4-5 million years ago."

"This means polar bears definitely persisted through warming periods during Earth's history," Lindqvist said. She cautions, however, that the species' endurance over several million years doesn't guarantee its future survival. To model historical populations of the polar bear, the scientists used computer simulations to analyze a deeply sequenced polar bear genome. "This is the first time we can see, from their genes, how the population history of polar bears tracked Earth's climate history," Lindqvist said. "We see an increase in polar bears at the end of the Early Pleistocene as the Earth became much colder, and a continuous decline in the size of the population during warmer times. We also found, perhaps unsurprisingly, that polar bears occur in much smaller numbers today than during prehistory," Lindqvist continued. "They have indeed lost a lot of their past genetic diversity, and because of this, they are very likely more sensitive to climate change threats today."

Discrepancies between the estimated age of polar bears in the new study and past studies could be explained by interbreeding between polar bears and brown bears since the species split from each other.

The new analysis uncovered more genetic similarities than previously known between polar bears and ABC brown bears, an isolated group from southeastern Alaska -- suggesting that these animals have exchanged genes since becoming separate species. "The ABC brown bears' mitochondrial sequences are much more like polar bears' than like other brown bears'," Miller said. "This made us wonder what other parts of their genomes are 'polar-bear-like'. We mapped such regions, which

constitute 5 to 10 percent of their total [DNA](#), onto the genomes of two ABC brown bears. As such, brown/polar bear hybridization, which has been observed recently in Arctic Canada, has undoubtedly contributed to shaping the modern polar bear's evolutionary story."

This intermingling between species is just one interesting finding emerging from the enormous trove of data that the PNAS study produced. Another question that the research is beginning to address: What makes a polar bear a polar bear? Polar bears have genetic differences from [brown bears](#) that let them survive in an Arctic climate with very different diets, and the new study identified genes that may be responsible for traits such as [polar bears'](#) pigmentation and the high fat content of their milk.

More information: "Polar and brown bear genomes reveal ancient admixture and demographic footprints of past climate change," by Webb Miller et al. *PNAS*, 2012.

Provided by Pennsylvania State University

Citation: Polar bear evolution tracked climate change, new DNA study suggests (2012, July 23) retrieved 18 April 2024 from

<https://phys.org/news/2012-07-polar-evolution-tracked-climate-dna.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.