

Scientists identify characteristics of successful malaria parasites in New Zealand bird populations

July 17 2012

Scientists from the Zoological Society of London (ZSL) are the first to document the characteristics of invading parasites, using malaria in New Zealand bird species.

The study, published today in <u>Ecology Letters</u>, identifies the factors influencing the success of parasites unintentionally introduced to new environments.

Avian malaria is a disease caused by species of parasites, of the genus *Plasmodium*, which infects birds. Just like human malaria, it is spread by mosquitoes, and the parasites spend part of their lives in <u>red blood cells</u> of birds. Avian malaria is common in continental areas, but is often absent from isolated islands where <u>mosquitoes</u> are less prevalent.

More than 800 exotic and native host birds were studied in a range of areas across Northern New Zealand. They detected <u>parasite infection</u> by extracting DNA from blood and analysing it to look at specific segments of genes. They then looked in more detail at the characteristics of the parasites they found to see if they had features that made them more likely to be present in bird hosts in New Zealand.

ZSL's Dr John Ewen, who is from New Zealand himself, says: "We have found a surprisingly high diversity of malaria parasites in New Zealand, including two found nowhere else in the world. However, most parasites



we found are recent arrivals, probably from infected birds released by humans. They tend to be widespread and common strains, which can infect a broad range of bird hosts. These findings will help us understand the what, when and how of exotic parasite introductions globally."

The global movement of parasites beyond their native country is an increasing problem, especially in the conservation of species. Many introduced parasites flourish in new environments and some can even be invasive. Little is known about the traits which enable parasites to survive and thrive.

ZSL's Professor Tim Blackburn added: "While we know a lot about the traits of successful vertebrate <u>species invasions</u>, our knowledge of the parasites they may carry is largely a blank page. This is alarming given that these co-introduced parasites can have catastrophic consequences for the natives they encounter, an example illustrated by smallpox introduced to the New World by the early European colonists."

This is the first study looking at the specific features of the exotic parasites within an isolated island bird community. New Zealand has a highly threatened bird community, sensitive to many human caused changes with exotic species being a major threat. ZSL scientists and colleagues are continuing to work on developing appropriate risk assessments for New Zealand's iconic native biodiversity and in understanding why some environments are more likely to be invaded by parasites than others.

Provided by Zoological Society of London

Citation: Scientists identify characteristics of successful malaria parasites in New Zealand bird populations (2012, July 17) retrieved 17 April 2024 from https://phys.org/news/2012-07-scientists-characteristics-successful-malaria-parasites.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.