

## Advance in X-ray imaging shines light on nanomaterials

August 7 2012

A new advance in X-ray imaging has revealed the dramatic threedimensional shape of gold nanocrystals, and is likely to shine a light on the structure of other nano-scale materials.

Described today in *Nature Communications*, the new technique improves the quality of nanomaterial <u>images</u>, made using X-ray diffraction, by accurately correcting distortions in the X-ray light.

Dr Jesse Clark, lead author of the study from the London Centre for Nanotechnology said: "With nanomaterials playing an increasingly important role in many applications, there is a real need to be able to obtain very high quality three dimensional images of these samples.

"Up until now we have been limited by the quality of our X-rays. Here we have demonstrated that with imperfect X-ray sources we can still obtain very high quality images of nanomaterials."

Up until now, most nanomaterial imaging has been done using <u>electron</u> <u>microscopy</u>. X-ray imaging is an attractive alternative as X-rays penetrate further into the material than electrons and can be used in ambient or controlled environments.

However, making lenses that focus X-rays is very difficult. As an alternative, scientists use the indirect method of coherent diffraction imaging (CDI), where the <u>diffraction pattern</u> of the sample is measured (without lenses) and inverted to an image by computer.



<u>Nobel Prize winner</u> Lawrence Bragg suggested this method in 1939 but had no way to determine the missing phases of the diffraction, which are today provided by <u>computer algorithms</u>.

CDI can be performed very well at the latest synchrotron X-ray sources such as the UK's Diamond Light Source which have much higher coherent flux than earlier machines. CDI is gaining momentum in the study of nanomaterials, but, until now, has suffered from poor image quality, with broken or non-uniform density. This had been attributed to imperfect coherence of the X-ray light used.

The dramatic <u>three-dimensional images</u> of gold <u>nanocrystals</u> presented in this study demonstrate that this distortion can be corrected by appropriate modelling of the coherence function.

Professor Ian Robinson, London Centre for Nanotechnology and author of the paper said: "The corrected images are far more interpretable that ever obtained previously and will likely lead to new understanding of structure of nanoscale materials."

The method should also work for free-electron-laser, electron- and atombased diffractive imaging.

**More information:** 'High-resolution three-dimensional partially coherent diffraction imaging' is published online in the journal *Nature Communications*.

## Provided by University College London

Citation: Advance in X-ray imaging shines light on nanomaterials (2012, August 7) retrieved 16 June 2024 from <u>https://phys.org/news/2012-08-advance-x-ray-imaging-nanomaterials.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.