


Thin current sheets in space: where the action is

August 1 2012



The four Cluster satellites in orbit round the Earth. (Artist's impression: ESA)


Much of the exciting action in space is confined to thin boundaries. The Universe is filled with plasma, a charged gas consisting of ions and electrons. Thin sheets with currents separate large plasma regions in space. Scientists at the Swedish Institute of Space Physics (IRF) have now finally measured the fundamental properties of one of the waves

mixing and accelerating plasmas within these sheets.

Around Earth, the processes accelerating electrons which hit the atmosphere and cause beautiful auroras are often initiated in thin current sheets. Similar processes, [auroras](#) and thin current sheets are found around other planets such as Jupiter and Saturn. Plasma regions close to the hot [solar surface](#) are separated by thin current sheets, and similar boundaries should also be common around [distant stars](#). In man-made plasmas, thin boundaries are found in the tokamak plasma employed in nuclear fusion research and [space observations](#) may help us understand fusion plasmas.

The [solar wind](#) blows plasma at the Earth's magnetic field. This causes the so-called magnetotail, stretching several hundred thousand kilometres downstream from the Earth. There is a thin current sheet separating the northern and southern parts of the tail.

In large parts of space, the plasma is too tenuous for the particles to actually collide. However, since the particles are charged, electric fields caused by some particles will interact with other particles. Often rather specific waves in the electric field interchange energy between the plasma particles. These waves replace ordinary collisions.



The electric field forms vortices propagating past two of the Cluster spacecraft. Since the electrons have much lower mass than the ions and can easily be moved around by the electric field, the electrons create a magnetic field, as shown in the bottom panel. (Fig. from Norgren et al., PRL, 2012)

The lower hybrid drift waves have been studied for 50 years and are thought to play an important role in these narrow current sheets. However, due to their relatively short wavelength, it has been impossible to observe their [fundamental properties](#). IRF's scientists have now, for the first time, been able to make direct measurements of the wavelength and velocity of these waves.

It has not been possible to measure the wavelength with a single spacecraft, but this can be done with the European Space Agency's four Cluster spacecraft. Taking advantage of the short 40 km separation between two of the four spacecraft in the magnetotail during August

2007, the scientists could observe the same wave propagating past first one and then the other spacecraft. The wavelength could be determined to be about 60 km (comparable to the radius of the electron gyro-motion in the magnetic field) and the velocity to about 1000 km/s (comparable to the ion velocity). The results appeared in the scientific journal *Physical Review Letters* on 31 July.

"We see small vortices that propagate in this narrow current sheet. They are just big enough so that both of the spacecraft can see them at the same time and be sure it is the same structure," says Cecilia Norgren of the Swedish Institute of [Space Physics](#) and a PhD student at Uppsala University. "The assumptions, used for several decades, have finally been verified by direct observations."

More information: Article in *Physical Review Letters*, vol. 109, issue 5 (2012): prl.aps.org/abstract/PRL/v109/i5/e055001

Provided by Swedish Institute of Space Physics

Citation: Thin current sheets in space: where the action is (2012, August 1) retrieved 3 May 2024 from <https://phys.org/news/2012-08-thin-current-sheets-space-action.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|