

For the first time, astronomers have measured the radius of a black hole

September 27 2012, by Jennifer Chu



This image, created using computer models, shows how the extreme gravity of the black hole in M87 distorts the appearance of the jet near the event horizon. Part of the radiation from the jet is bent by gravity into a ring that is known as the 'shadow' of the black hole. Credit: Avery E. Broderick (Perimeter Institute & University of Waterloo

(Phys.org)—The point of no return: In astronomy, it's known as a black



hole—a region in space where the pull of gravity is so strong that nothing, not even light, can escape. Black holes that can be billions of times more massive than our sun may reside at the heart of most galaxies. Such supermassive black holes are so powerful that activity at their boundaries can ripple throughout their host galaxies.

Now, an international team, led by researchers at MIT's Haystack Observatory, has for the first time measured the radius of a black hole at the center of a <u>distant galaxy</u>—the closest distance at which matter can approach before being irretrievably pulled into the black hole.

The scientists linked together <u>radio dishes</u> in Hawaii, Arizona and California to create a <u>telescope array</u> called the "<u>Event Horizon</u> Telescope" (EHT) that can see details 2,000 times finer than what's visible to the <u>Hubble Space Telescope</u>. These radio dishes were trained on M87, a galaxy some 50 million light years from the Milky Way. M87 harbors a black hole 6 billion times more massive than our sun; using this array, the team observed the glow of matter near the edge of this black hole—a region known as the "event horizon."

"Once objects fall through the event horizon, they're lost forever," says Shep Doeleman, assistant director at the MIT Haystack Observatory and research associate at the Smithsonian Astrophysical Observatory. "It's an exit door from our universe. You walk through that door, you're not coming back."

Doeleman and his colleagues have published the results of their study this week in the journal *Science*.

Jets at the edge of a black hole

Supermassive black holes are the most extreme objects predicted by <u>Albert Einstein's theory of gravity</u>—where, according to Doeleman,



"gravity completely goes haywire and crushes an enormous mass into an incredibly close space." At the edge of a black hole, the <u>gravitational</u> <u>force</u> is so strong that it pulls in everything from its surroundings. However, not everything can cross the event horizon to squeeze into a black hole. The result is a "cosmic traffic jam" in which gas and dust build up, creating a flat pancake of matter known as an <u>accretion disk</u>. This disk of matter orbits the black hole at nearly the speed of light, feeding the black hole a steady diet of superheated material. Over time, this disk can cause the black hole to spin in the same direction as the orbiting material.



An accretion disk (orange) of gas and dust surrounds super-massive black holes at the center of most galaxies. These disks of galactic matter emit magnetic



beams (pink lines) that spew out from the center of the black hole, drawing matter out from both ends in high-powered jets. Credit: NASA and Ann Field

Caught up in this spiraling flow are magnetic fields, which accelerate hot material along powerful beams above the accretion disk The resulting high-speed jet, launched by the black hole and the disk, shoots out across the galaxy, extending for hundreds of thousands of light-years. These jets can influence many galactic processes, including how fast stars form.

'Is Einstein right?'

A jet's trajectory may help scientists understand the dynamics of black holes in the region where their gravity is the dominant force. Doeleman says such an extreme environment is perfect for confirming Einstein's theory of general relativity—today's definitive description of gravitation.

"Einstein's theories have been verified in low-gravitational field cases, like on Earth or in the solar system," Doeleman says. "But they have not been verified precisely in the only place in the universe where Einstein's theories might break down—which is right at the edge of a black hole."

According to Einstein's theory, a black hole's mass and its spin determine how closely material can orbit before becoming unstable and falling in toward the event horizon. Because M87's jet is magnetically launched from this smallest orbit, astronomers can estimate the black hole's spin through careful measurement of the jet's size as it leaves the black hole. Until now, no telescope has had the magnifying power required for this kind of observation.

"We are now in a position to ask the question, 'Is Einstein right?""



Doeleman says. "We can identify features and signatures predicted by his theories, in this very strong gravitational field."



This artist's impression of the innermost regions of M87 shows the relationship between the black hole, the orbiting accretion flow, and the launching of the relativistic jet. Credit: Perimeter Institute for Theoretical Physics

The team used a technique called Very Long Baseline Interferometry, or VLBI, which links data from radio dishes located thousands of miles



apart. Signals from the various dishes, taken together, create a "virtual telescope" with the resolving power of a single telescope as big as the space between the disparate dishes. The technique enables scientists to view extremely precise details in faraway galaxies.

Using the technique, Doeleman and his team measured the innermost orbit of the accretion disk to be only 5.5 times the size of the black hole event horizon. According to the laws of physics, this size suggests that the accretion disk is spinning in the same direction as the black hole—the first direct observation to confirm theories of how black holes power jets from the centers of galaxies.

The team plans to expand its telescope array, adding radio dishes in Chile, Europe, Mexico, Greenland and Antarctica, in order to obtain even more detailed pictures of <u>black holes</u> in the future.

More information: www.eventhorizontelescope.org/

Ref. "Jet Launching Structure Resolved Near the Supermassive Black Hole in M87," by S.S. Doeleman et al., *Science*, 2012.

Provided by Massachusetts Institute of Technology

Citation: For the first time, astronomers have measured the radius of a black hole (2012, September 27) retrieved 1 May 2024 from <u>https://phys.org/news/2012-09-astronomers-radius-black-hole.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.