


Within reach of Australia: Rabies is now present only 350 kilometres from northern Australia

September 28 2012



Rabies, spread mainly by rabid dogs, kills an estimated 100,000 people a year.
Credit: Flickr/Andy Wagstaffe

Rabies is now present only 350 kilometres from northern Australia, closer than the distance from Sydney to Dubbo.

If it reached our shores it would have serious ecological, public health, economic and social impacts according to Professor Michael Ward, from the University of Sydney.

Professor Ward is conducting research to identify how [rabies](#) spreads

and the risks of future outbreaks, such as evaluating [surveillance systems](#) to detect a potential incursion in the Top End and Torres Strait.

Speaking on World Rabies Day, 28 September, Professor Ward, the Chair of Veterinary Public Health and Food Safety at the University of Sydney said, "Rabies, spread mainly by the bite of rabid dogs, and inevitably fatal, is a serious health risk for communities in Asia and Africa. The disease kills an estimated 100,000 people a year.

"Australia is fortunate to not be affected by this terrible disease, which infects the [central nervous system](#) and causes inflammation of the brain. However, during the past 10 to 15 years rabies has spread to areas of [eastern Indonesia](#) that were previously rabies-free, such as Flores and Bali. In 2010, the disease reached the Tanimbar Islands, part of the Moluccas, just 350 kilometres north of the Top End."

The University's Faculty of Veterinary Science is researching rabies in Indonesia and its potential to spread to neighbouring regions that are currently free of the disease, specifically Timor Leste, [Papua New Guinea](#) and [Northern Australia](#).

"Our work overseas involves observing and recording when and how often dogs are being transported between islands including direct observations of ferries and [fishing boats](#). Estimating the size of the dog population in rabies-free areas is also an important component of determining risk," said Professor Ward.

In northern Australia, the research team from the faculty is applying a similar approach, recording the movement of dogs between communities to determine how rabies might spread, should it be introduced.

While some government surveillance for rabies is already being undertaken the team is assessing additional surveillance systems to better

detect a rabies incursion.

"We are concerned about the potential ecological impact of rabies in this region," Professor Ward said.

"Dingoes, as a top predator, play a crucial role in maintaining ecosystem diversity and if their numbers were substantially reduced, it could affect many other species in this fragile ecosystem.

"Potential economic impacts on the cattle industry should also not be underestimated. Cattle losses from rabies can be significant and would be very hard to prevent if rabies got into the Top End. And of course, controlling rabies would be horrendously expensive."

"To prevent rabies entering Australia, we must help all our near neighbours to increase their alertness and preparedness, and also help Indonesia to reverse this seemingly inexorable spread of rabies to the southeast," said Dr Helen Scott-Orr, an affiliate of the Faculty of [Veterinary Science](#), current board director of the Invasive Animals Cooperative Research Centre and Animal Health Australia, and former Chief Veterinary Officer of NSW.

"It is really unacceptable that, in this day and age, a dreaded disease like rabies should continue to ravage new vulnerable communities near us."

Experience in Flores, Bali and elsewhere shows that to eradicate rabies we must vaccinate dogs against the disease and maintain a level of 70 percent vaccination coverage across the dog population, both tame and wild, for several years, Dr Scott-Orr says.

"This sounds simple but is extraordinarily difficult and also very costly," said Dr Scott-Orr. "Culling wild or stray dogs, which may seem logical, just doesn't work, and can actually do more harm than good."

Provided by University of Sydney

Citation: Within reach of Australia: Rabies is now present only 350 kilometres from northern Australia (2012, September 28) retrieved 18 April 2024 from <https://phys.org/news/2012-09-australia-rabies-kilometres-northern.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.