

Glacial youth therapy for the Scandinavian landscape

September 24 2012, by Sverre Ole Drønen



The fjords of Norway are on UNESCO's World Heritage List and are often considered as the most scenically outstanding fjords of the planet. Foto: Colourbox

The high elevation flat surfaces characteristic of the Norwegian landscape are in geologically terms young, according to a paper in *Nature Geoscience*.

In a paper recently published in *Nature [Geoscience](#)*, researchers from the University of Bergen (UiB) and ETH Zurich have demonstrated that ice sheets have extensively shaped the fjords of Norway for the last 2.8 million years.

"However, the massive [sediment record](#) deposited offshore Norway during this period tells us a more complex story," explains Postdoctoral

Fellow Philippe Steer at UiB's Department of Earth Science and the first author of the paper. "Ice sliding during [glacial periods](#) also resulted in eroding the present-day flattish surfaces that form the high-elevation parts of the Scandinavian landscape."

Contradicting previous views

The conclusion of this new study by Steer and his collaborators in Bergen and Zurich contradicts previously accepted views for Scandinavian geodynamics, where high-elevation parts of the landscape were left almost untouched by the repeated ice sheets.

"The origin and fate of the western Scandinavian landscapes under the repeated ice sheets of the last 2.8 million years has been a highly debated scientific topic for a hundred years," says Professor Ritske S. Huisman, head of the research project and a co-author of the paper. "In our contribution we have, for the first time, managed to demonstrate that the [glaciers](#) and ice sheets have eroded away both the Scandinavian fjords and flat high-elevation surfaces during the past glacial times. This supports the idea that the mountain range in western Scandinavian as seen today is a remnant of the more than 400 million years old Caledonian mountain range and has been at high elevations ever since."

The researchers believe that this new conceptual model of topographic glacial imprint may also apply to other high-latitude landscapes, such as the one in Greenland.

"This discovery was made possible by linking two fields, onshore and offshore geology that are usually studied separately. Because onshore erosion results ultimately in offshore sediments, it is fundamental to understand the past sediment record to assess the history of Scandinavian landscapes," Huisman explains. "Our research indicates that erosion must also have occurred in the areas above the fjords, e.g. on top of the

flat surfaces and demonstrates that these high elevation flat surfaces characteristic of the Norwegian landscape are in geological terms young."

The study was made possible because of the large amount of data available for research—such as seismic profiles and drill cores—that was acquired offshore Norway by Statoil and other companies. This data offers a very precise image of the Scandinavian sediment history.

Reshaped by glacial erosion

By demonstrating that the entire topography was reshaped by glacial erosion, the study asks for a complete re-interpretation of the tectonic history of Scandinavia.

"This result has important consequences for oil and gas exploration offshore Norway as it aids in the reconstruction of topography in the past, and may also help to better understand past topographic and tectonic evolution of other less-well constrained glaciated areas, such as Greenland. In addition, this study offers a new insight into the landscapes and fjords of western Scandinavia, which are considered to be among the most scenically outstanding areas on the planet, the Nærøyfjord and Geirangerfjord being on the UNESCO world heritage list," Steer says.

More information: Bimodal Plio–Quaternary glacial erosion of fjords and low-relief surfaces in Scandinavia. Philippe Steer, Ritske S. Huisman, Pierre G. Valla, Sébastien Gac & Frédéric Herman. *Nature Geoscience*. Volume: 5, Pages: 635–639. Year published: (2012). DOI: [doi:10.1038/ngeo1549](https://doi.org/10.1038/ngeo1549)

Provided by University of Bergen

Citation: Glacial youth therapy for the Scandinavian landscape (2012, September 24) retrieved 4 May 2024 from <https://phys.org/news/2012-09-glacial-youth-therapy-scandinavian-landscape.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.