

# Malay Archipelago bat not one, but two species

September 7 2012

---



*Myotis muricola* from Kuningan, West Java. Otherwise known as the Wall-roosting Mouse-eared bat, it is a species of vesper bat. They are typically insectivorous and nocturnal.

Genetic studies of *Myotis muricola*, otherwise known as the Wall-roosting Mouse-eared bat or Nepalese Whiskered *Myotis*, suggest that it consists of not one, but two distinct species.

*M. muricola* is widespread on the Malay Archipelago, a region with an island geography that provides natural boundaries and as a result displays some of the richest biodiversity in the world. Until now scientists had compared the shape and size of *Myotis* bats across the Archipelago and conducted only limited [DNA analysis](#). As a result the *Myotis* family tree has remained in disarray with often contradictory lines of evidence confusing matters.




Fig. 2 The localities of the specimens of *M. muricola* that were examined. The west of Wallace's Line region is represented by (1) Jambi (2) Krakatau Island (3) Bali Island (4) East Kalimantan (5) Southern Sarawak (6) Central Sarawak (7) Northern Sarawak (8) Tawau, Sabah (9) Fraser Hill, Pahang. The sites from the east of Wallace's Line include: (10) Lombok Island (11) Sumbawa Island (12) Sumba Island (13) Flores Island (14) Lembata Island (15) Pantar Island (16) Timor Island

To resolve these taxonomic difficulties, a team from the Universiti Malaysia Sarawak and the Indonesian Institute of Sciences launched the first in-depth genetic study on *M. muricola* and its sister taxa, *M. mystacinus*. Writing in the *Pertanika Journal of Tropical Agricultural Science*, the team describe how they examined the bats in two regions, West and East of Wallace's line (fig. 2). Using a [genetic analysis](#) technique known as 'DNA cytochrome b sequencing' (fig. 3), they found that the Western and Eastern groups had a genetic distance of between

26 and 39 per cent.

Based on this high [genetic distance](#) they argue that *M. muricola* Eastern and *M. muricola* Western should be considered as two [distinct species](#). The data also suggest that the Eastern bat started to diversify in the western region during the Pliocene (5 to 2.5 million years ago). They became fully diverged within the western region during the Pleistocene (2.5 million to 11,000 years ago) under the influence of ancient Sunda River systems that had produced gallery forest corridors which functioned as safe havens for the bats as the climate and geography around them changed.




Fig. 3 Cytochrome b is a component of mitochondrial DNA (mtDNA) which is frequently used to determine the evolutionary relationships between organisms as it is highly variable and therefore useful for determining relationships within families and genera.

The authors stress that as a result of these findings an official revision of the taxonomic status of *M. muricola* is urgently needed. In addition, more samples from throughout the geographic range are required to

firmly establish these findings.

**More information:** [www.pertanika.upm.edu.my/Pertanika%20PAPERS/JTAS%20Vol.%2035%20%282%29%20May.%202012/12%20Pg%20271-292.pdf](http://www.pertanika.upm.edu.my/Pertanika%20PAPERS/JTAS%20Vol.%2035%20%282%29%20May.%202012/12%20Pg%20271-292.pdf)

Provided by Universiti Putra Malaysia (UPM)

Citation: Malay Archipelago bat not one, but two species (2012, September 7) retrieved 8 May 2024 from <https://phys.org/news/2012-09-malay-archipelago-species.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--