


X-rays reveal the self-defence mechanisms of bacteria

September 14 2012



The toxins normally bind very strongly to the antitoxins and are thus not only inactive, but also prevent the production of more toxin from the information encoded in the bacterial DNA. During the dormant state, however, the antitoxins are degraded, and the toxins released (step 1). The free toxins now bind to unoccupied antitoxins on DNA within the area encoding the toxin-antitoxin couple (step 2). Binding increasing amounts of toxin eventually leads to the release of the molecules from the gene (steps 3 and 4) and finally to new toxin production. Credit: Ditlev E. Brodersen

A research group at Aarhus University has gained unique insight into how bacteria control the amount of toxin in their cells. The new findings can eventually lead to the development of novel forms of treatment for bacterial infections.

Many pathogenic bacteria are able to go into a dormant state by producing persister cells that are not susceptible to conventional antibiotics. This causes serious problems in the treatment of life-threatening diseases such as tuberculosis, where the presence of persister

cells often leads to a resurgence of infection following [medical treatment](#)

At the molecular level, the formation of persister cells is due to the presence of toxins that are produced by the bacteria themselves, and which enable them to enter the dormant state. During this hibernation period, the bacteria constantly regulate the amount of toxin at exactly the same level and thus maintain the dormant state.

In an article recently published in the American scientific journal *Structure*, the researchers at the Department of [Molecular Biology](#) and Genetics, Aarhus University, present new results that reveal the molecular details of the [regulatory mechanism](#) of toxins.

By isolating and crystallising the toxin molecules and their molecular companions – the antitoxins – and by subsequently exposing the crystals to strong X-rays, the research team (consisting of the two PhD students Andreas Bøggild and Nicholas Sofos and Associate Professor Ditlev E. Brodersen) gained unique insight into how bacteria control the amount of toxin in the cell (illustrated in the info box).

The new findings can eventually lead to the development of entirely new forms of treatment of bacterial infections that work initially by blocking [toxin](#) function and production, and subsequently by using traditional antibiotics to fight the [pathogenic bacteria](#).

More information: [www.sciencedirect.com/science/ ...
ii/S0969212612002997](http://www.sciencedirect.com/science/.../S0969212612002997)

Provided by Aarhus University

Citation: X-rays reveal the self-defence mechanisms of bacteria (2012, September 14) retrieved 11 May 2024 from <https://phys.org/news/2012-09-x-rays-reveal-self-defence-mechanisms-bacteria.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.