

## Research aims to extend strawberry growing season in Mid-Atlantic region

October 25 2012, by Sharon Durham

---



Reaching inside a low tunnel that extends the fruit's growing season, geneticist Kim Lewers and horticulturalist John Enns observe strawberries with anthracnose fruit rot. The next goal is to transfer disease resistance to future varieties to reduce disease incidence. Credit: Stephen Ausmus

Growing strawberry plants in the U.S. Mid-Atlantic region beneath canopy-like structures called low tunnels can allow the season to start earlier and continue through the summer and fall, according to U.S. Department of Agriculture (USDA) scientists.

At the Agricultural Research Service (ARS) [Genetic Improvement of Fruits and Vegetables](#) Laboratory in Beltsville, Md., geneticist Kim Lewers is testing some strawberry cultivars in the new production system designed to extend the growing season in the northern and eastern United States. Lewers' research partners are horticulturist John Enns and support services staffer George Meyers. ARS is USDA's chief intramural scientific research agency.

Low tunnels are canopies made of long sheets of plastic laid over support hoops that hold it about 30 inches above the strawberry bed. Strawberries are planted beneath these structures, which protect the fruit from rain, provide shade from damaging infrared and UV light, and can capture warmth during the cooler spring and fall seasons.

By protecting the plants from rain, the tunnels help to minimize two important diseases of strawberry, Botrytis and anthracnose, which thrive in the rain. Botrytis occurs in cool, [wet conditions](#), while anthracnose takes hold in hot, wet conditions.

According to Lewers, high tunnels also are used by growers, but they can be problematic because the humidity is higher in the tunnel, which causes more Botrytis and more [powdery mildew](#), another strawberry disease. But in low tunnels, the humidity is the same as outside the tunnel when the sides of the tunnel are up.

The Maryland peak [strawberry](#) season is usually mid-May to mid-June, but in Lewers' low-tunnel production system, strawberries start earlier and continue through the fall—essentially a whole new season. In some months, yields in the low tunnels can be as high as those from the same cultivars when they are grown in California, where they were developed.

All of the [strawberry plant](#) material developed in Lewers' research program has been and still is freely available. The plants are not

patented, so they are available without special license to any nursery that wants to grow them.

Read more about this and other fruit and vegetable research in the October 2012 issue of *Agricultural Research* magazine.

**More information:** [www.ars.usda.gov/is/AR/archive ...  
oct12/fruits1012.htm](http://www.ars.usda.gov/is/AR/archive...oct12/fruits1012.htm)

Provided by Agricultural Research Service

Citation: Research aims to extend strawberry growing season in Mid-Atlantic region (2012, October 25) retrieved 27 April 2024 from <https://phys.org/news/2012-10-aims-strawberry-season-mid-atlantic-region.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.