

Genome evolution and carbon dioxide dynamics

October 25 2012, by Dr. Ilia Leitch



Impact of genome size on guard cell size. *Paris japonica* (A) and *Quercus robur* (B) differ 170-fold in genome size and this is reflected in the size of the guard cells. Both pictures are taken at the same magnification. Credit: Peter Franks

Using the size of guard cells in fossil plants to predict how much DNA each cell contained (the genome size), researchers have discovered that variations in genome sizes over geological time correlate with atmospheric carbon dioxide levels.

The total amount of DNA in the nucleus of an organism is known as the genome size. In plants this character varies nearly 2400-fold between

species, with many having up to 50 times more DNA than is found in our own genome.

Fossil guard cells reveal genome size evolution

Understanding the biological significance of this variation has been an ongoing research theme at Kew for nearly 30 years. One area of focus has been the study of genome size evolution and great progress has been made by superimposing genome size data onto the phylogenetic trees of plants. Nevertheless, we have had no idea of how genome size has varied over geological time as it is impossible to measure this from fossils.

Given that the size of genomes and [guard cells](#) (which form the [stomata](#)) are closely linked (see picture), researchers at the universities of Sheffield (UK) and Yale (USA), in collaboration with Kew scientists, used the size of fossil guard cells as a proxy for genome size to track how this has changed over the past 400 million years (spanning the full geological history of vascular plants). The results suggest that genome size evolution has been dynamic with both increases and decreases taking place during [plant evolution](#).

Genome sizes correlate with carbon dioxide levels

Coupling this information with knowledge of how atmospheric carbon dioxide levels have fluctuated over the same time period revealed a good correlation between these parameters, with larger guard cells and inferred genome sizes found when levels of carbon dioxide were high. Of course, the observed correlation does not mean that carbon dioxide levels are directly driving changes in guard cell size and genome size. Nevertheless, because guard cells play a key role in optimising gas exchange in plants, the results hint at the possibility that carbon dioxide might be one of the selective forces responsible for generating the

diversity of guard cell and genome sizes encountered in today's flora.

Given that the particular genome size of a plant can have a considerable influence on where, when and how a plant lives, understanding the nature of the relationship between carbon dioxide levels and [genome size](#) will become increasingly important as the level of [atmospheric carbon dioxide](#) continues to rise.

More information: Franks, P. J., Leitch, I. J., Ruszala, E. M., Hetherington, A. M. & Beerling, D. J. (2012). Physiological framework for adaptation of stomata to CO₂ from glacial to future concentrations. *Philosophical Transactions of the Royal Society B* 367: 537-546.
Franks, P. J., Freckleton, R. P., Beaulieu, J. M., Leitch, I. J. & Beerling, D. L. (2012). Megacycles of atmospheric carbon dioxide concentration correlate with fossil plant genome size. *Philosophical Transactions of the Royal Society B* 367: 556-564

Provided by Royal Botanic Gardens, Kew

Citation: Genome evolution and carbon dioxide dynamics (2012, October 25) retrieved 23 May 2024 from <https://phys.org/news/2012-10-genome-evolution-carbon-dioxide-dynamics.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--