

Testing Mars and Moon soil for sheltering astronauts from radiation

October 10 2012



NASA proposal for a future deep space station, 'parked' at lunar Lagrange point 2 past the far side of the Moon. Residing beyond Earth's magnetic field, its occupants would need enhanced radiation protection. Credits: NASA



(Phys.org)—Humans venturing beyond Earth orbit deeper into space face increased exposure to cosmic radiation, so ESA has teamed with Germany's GSI particle accelerator to test potential shielding for astronauts, including Moon and Mars soil.

ESA's two-year project is assessing the most promising materials for shielding future astronauts going to the Moon, the asteroids or Mars.

"We are working with the only facility in Europe capable of simulating the high-energy heavy atomic nuclei found in galactic cosmic radiation – the GSI Helmholtz Centre for <u>Heavy Ion Research</u> in Darmstadt, Germany," explained Alessandra Menicucci, overseeing the project.

"We assessed materials including aluminium, water, polyethylene plastic, multilayer structures and simulated Moon and Mars material – the latter on the basis these will be accessible to planetary expeditions.

"We have also confirmed a new type of <u>hydrogen storage</u> material holds particular promise."

Space is awash with charged particles, meaning that astronauts are officially classed as radiation workers.





Lunar regolith simulant JSC-1A Credits: NASA

The <u>International Space Station</u> orbits within Earth's magnetic field, safeguarding its occupants from the bulk of space radiation. To venture further out, dedicated shielding will be required.

<u>Space radiation</u> comes from the Sun – in the form of intense but shortlived 'solar particle events' – as well as galactic cosmic radiation originating beyond our Solar System: atomic nuclei produced by dying stars, their passage sped by magnetic fields as they cross the galaxy.

"Solar particle events are made up of protons that can be shielded quite simply," added Alessandra.

"The real challenge for deep-space missions is galactic cosmic radiation, which cannot be shielded completely because of its very high energy, although the exposure level decreases with increased solar activity.



Most are small protons or helium nuclei, but about 1% are larger, the size of an iron atom or more – known as 'high-ionising high energy particles' or HZE for short.



GSI's linear accelerator UNILAC (UNIversal Linear ACcelerator) has a length of 120 meters. Ions, e.g. charged atoms of all chemical elements can be accelerated up to 20 percent of the speed of light (60,000 km/s) with this unit. Credits: GSI



Radiation shielding can be counter-intuitive because denser and thicker does not always mean better.

HZEs striking metal shields can produce showers of secondary particles that might be even more harmful.

And as shield thickness increases, overall the energy loss of ionising radiation rises to a peak then declines rapidly.

"In general, the lighter a material's <u>atomic nuclei</u> the better the protection," notes Alessandra.

Water and polyethylene performed better than aluminium for instance, and new hydrogen-rich materials developed by UK company Cella Energy tested better still.

Cella Energy originally developed its patent-pending materials for storing hydrogen fuel but is currently investigating their radiation resistance.

Provided by European Space Agency

Citation: Testing Mars and Moon soil for sheltering astronauts from radiation (2012, October 10) retrieved 3 May 2024 from <u>https://phys.org/news/2012-10-mars-moon-soil-astronauts.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.