

Interstellar dust and the sun

November 12 2012



An artist's concept of the heliosphere (seen in blue, including a shocked region). The Earth is at 1 AU, and the two Voyager spacecraft are seen beyond 100 AU (the Cassini spacecraft at Saturn is also shown). A new study investigates what happens to interstellar dust that encounters the solar system and the Sun's heliosphere. Credit: NASA and JHU/APL

(Phys.org)—The space between stars is not empty. It contains copious but diffuse amounts of gas and dust; in fact about 5-10% of the total mass of our Milky Way galaxy is in interstellar gas. About 1% of the



mass of this interstellar material, quite a lot in astronomical terms, is in the form of tiny dust grains made predominantly of silicates (sand too is made of silicates), though some grains are also composed of carbon and other elements. Dust grains are important. They block visible light while emitting infrared light, and thus help determine what astronomers can see while controlling much of the energy balance in the interstellar medium (ISM) by virtue of the absorption and subsequent re-emission at longer wavelengths of light from stars. Dust is also essential to the chemistry that takes place in the ISM because it provides gas molecules with a surface on which to react with other molecules. Not least, dust contains a large fraction of many important elements in the universe like silicon, carbon, and iron. Moreover, astronomers think that at some stage in the evolution of new stars the dust around them will coagulate into large clumps—the first step towards forming planets.

CfA astronomer Jonathan Slavin and a team of six other astronomers wondered what happens to interstellar dust when it wanders into the solar system and gets close enough to the Sun to fall under the influence of its radiation, winds, and gravity. They note that the Sun (and its planets) is moving through a low density cloud of partially ionized gas. This motion, together with the wind of particles that the Sun emits, produces a bowshaped region called the heliosphere, the bow-shaped end of which is about 100 AU from the Sun (one AU is the average distance of the Earth from the Sun).

Writing in the latest issues of the <u>Astrophysical Journal</u>, the scientists report on the results of their theoretical models of the behavior of interstellar dust grains as the Sun moves through space. They build on insitu observations of the heliosphere taken when the Voyager 1 and Voyager 2 spacecraft on their outward journey encountered the edges of the heliosphere, results that constrain its size and shape. Assuming typical grains made of olivine silicates, the team finds that the small grains (less than the wavelength of ultraviolet light) stay far away from



the Sun, that gravity helps the large grains collect near the Sun, but that intermediate-sized grains - about the size of the wavelength of optical light - can actually pile up in diffuse structures at the edges of the heliosphere. The new results, besides providing important new information on dust grains in the solar system, suggest that radiation from these intermediate-sized grain structures could contaminate the images of the sky used to measure the cosmic backgrounds.

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: Interstellar dust and the sun (2012, November 12) retrieved 26 April 2024 from <u>https://phys.org/news/2012-11-interstellar-sun.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.