

# What's eating the sea butterfly? CO<sub>2</sub>, study says

November 25 2012

---



Shore of Deception Island in Antarctica, pictured in 2008. Rising acidity is eating away the shells of tiny snails, known as "sea butterflies", that live in the seas around Antarctica, leaving them vulnerable to predators and disease, scientists said Sunday.

Rising acidity is eating away the shells of tiny snails, known as "sea butterflies", that live in the seas around Antarctica, leaving them vulnerable to predators and disease, scientists said Sunday.

The study presents rare evidence of living creatures suffering the results of ocean acidification caused by rising carbon dioxide levels from fossil fuel burning, the [British Antarctic Survey](#) said in a statement.

"The finding supports predictions that the impact of ocean acidification on [marine ecosystems](#) and food webs may be significant."

The tiny snail, named for two wing-like appendices, does not necessarily die as a result of losing its shell, but it becomes an easier target for fish and bird predators, as well as infection.

This may have a follow-through effect on other parts of the food chain, of which they form a core element.

The world's oceans absorb more than a quarter of man-made [carbon dioxide emissions](#), which lower the sea water pH.

Since the beginning of the industrial era, our oceans have become 30 percent more acidic, reaching an acidity peak not seen in at least 55 million years, scientists say.

Scientists discovered the effects of acidification on the sea butterflies from samples taken around the Scotia Sea region of the Southern Ocean in February 2008.

**More information:** Extensive dissolution of live pteropods in the Southern Ocean, *Nature Geoscience* (2012) [doi:10.1038/ngeo1635](https://doi.org/10.1038/ngeo1635)

## **Abstract**

The carbonate chemistry of the surface ocean is rapidly changing with ocean acidification, a result of human activities. In the upper layers of the Southern Ocean, aragonite—a metastable form of calcium carbonate with rapid dissolution kinetics—may become undersaturated by 2050.

Aragonite undersaturation is likely to affect aragonite-shelled organisms, which can dominate surface water communities in polar regions. Here we present analyses of specimens of the pteropod *Limacina helicina antarctica* that were extracted live from the Southern Ocean early in 2008. We sampled from the top 200 m of the water column, where aragonite saturation levels were around 1, as upwelled deep water is mixed with surface water containing anthropogenic CO<sub>2</sub>. Comparing the shell structure with samples from aragonite-supersaturated regions elsewhere under a scanning electron microscope, we found severe levels of shell dissolution in the undersaturated region alone. According to laboratory incubations of intact samples with a range of aragonite saturation levels, eight days of incubation in aragonite saturation levels of 0.94–1.12 produces equivalent levels of dissolution. As deep-water upwelling and CO<sub>2</sub> absorption by surface waters is likely to increase as a result of human activities, we conclude that upper ocean regions where aragonite-shelled organisms are affected by dissolution are likely to expand.

[Press release](#)

(c) 2012 AFP

Citation: What's eating the sea butterfly? CO<sub>2</sub>, study says (2012, November 25) retrieved 25 April 2024 from <https://phys.org/news/2012-11-sea-butterfly-co2.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---